



## PRINCIPLES AND MODELS OF CARE

### POST ACQUIRED BRAIN INJURY

Nora Cullen MSc MD FRCPC, Matthew J Meyer PhD,  
Heather MacKenzie MD FRCPC, Jo-Anne Aubut BA,  
Mark Bayley MSc MD FRCPC, Robert Teasell MD FRCPC

## Disclaimer

This review has been prepared based on the scientific and professional information available up to March 2020. The ERABI information is provided for informational and educational purposes only. If you have or suspect you have a health problem, you should consult your health care provider. The ERABI contributors shall not be liable for any damages, claims, liabilities, costs, or obligations arising from the use or misuse of this material.

## Copyright

With the exception of those portions of this document for which a specific prohibition or limitation against copying appears, the balance of this document may be reproduced and published in its entirety, without modification, in any form, including in electronic form, for educational or non-commercial purposes. Should any adaptation of the material be required for any reason, written permission must be obtained from ERABI. Appropriate credit or citation must appear on all copied material as follows:

Cullen N, Meyer M, MacKenzie H, Aubut J, Bayley M, Teasell R. (2019). Principles and Models of Care Following an Acquired Brain Injury. In Teasell R, Cullen N, Marshall S, Janzen S, Faltynek P, Bayley M, editors. Evidence-Based Review of Moderate to Severe Acquired Brain Injury. Version 13.0: p1-65.

## Funding

This work is funded by the Ontario Neurotrauma Foundation. All work produced by ERABI is editorially independent from its funding source.

## Conflict of Interest

In the context of ERABI development, the term “conflict of interest” (COI) refers to situations in which an author or ERABI staff member’s financial, professional, intellectual, personal, organizational or other relationships may compromise their ability to independently conduct this evidence-based review. No limiting conflicts were identified.

## Contact Information

### **Evidence-Based Review of Moderate to Severe Acquired Brain Injury**

550 Wellington Rd South, London, Ontario, Canada N6C 0A7

Website: [www.ERABI.ca](http://www.ERABI.ca)

# Greetings from Dr. Robert Teasell,

Professor and Chair-Chief of Physical Medicine and Rehabilitation



The Collaboration of Rehabilitation Research Evidence (CORRE) team is delighted to present the Evidence-Based Review of moderate to severe Acquired Brain Injury (ERABI) *Mental Health Issues post Acquired Brain Injury*. Through collaboration of researchers, clinicians, administrators, and funding agencies, ERABI provides an up-to-date review of the current evidence in brain injury rehabilitation. ERABI synthesizes the research literature into a utilizable format, laying the foundation for effective knowledge transfer to improve healthcare programs and services.

We offer our heartfelt thanks to the many stakeholders who are able to make our vision a reality. Firstly, we would like to thank the Ontario Neurotrauma Foundation, which recognizes ERABI's capacity to lead in the field of brain injury evidence-based reviews and is committed to funding it. We would also like to thank the co-chairs of ERABI, Dr. Mark Bayley (University of Toronto) and Dr. Shawn Marshall (University of Ottawa) for their invaluable expertise and stewardship of this review. Special thanks to the authors for generously providing their time, knowledge and perspectives to deliver a rigorous and robust review that will guide research, education and practice for a variety of healthcare professionals. We couldn't have done it without you! Together, we are building a culture of evidence-based practice that benefits everyone.

We invite you to share this evidence-based review with your colleagues, patient advisors that are partnering within organizations, and with the government agencies with which you work. We have much to learn from one another. Together, we must ensure that patients with brain injuries receive the best possible care every time they require rehabilitative care – making them the real winners of this great effort!

Robert Teasell, MD FRCPC

# Table of Contents

**Preface..... 5**  
    About the Authors .....5  
    Purpose .....6  
    Key Concepts.....6  
    Methods .....8  
    Interpretation of the Evidence .....9  
    Strength of the Evidence .....10

**Summary of the Evidence ..... 12**

**Introduction ..... 16**

**Acute Management and Implementation of Care ..... 19**  
    Implementation of Guidelines .....20  
    Institutional Comparisons.....26

**Inpatient Rehabilitation..... 28**  
    Intensity of Inpatient Rehabilitation .....31  
    Timing of Inpatient Rehabilitation .....36

**Outpatient Rehabilitation and Community Reintegration..... 40**  
    Vocational Rehabilitation .....47

**Complete Care Pathways ..... 49**

**Conclusion ..... 52**

**Reference List..... 54**


# Preface

## About the Authors

ERABI is internationally recognized and led by a team of clinicians and researchers with the goal of improving patient outcomes through research evidence. Each ERABI module is developed through the collaboration of many healthcare professionals and researchers.




Dr. Cullen is the newly appointed Division Director for PM&R in the Department of Medicine and Chief of PM&R at HHSC and St Joseph’s Healthcare. She has been the Chief of Staff and Chair of the Medical Advisory Committee at West Park Healthcare Centre for the last 10 years, actively seeking to promote system change and patient advocacy in an integrated system. Her goal is to enhance the quality of life of people with disabilities based on current best evidence.



Matthew J Meyer PhD



Heather MacKenzie, MD, FRCPC is a consultant physiatrist in the spinal cord injury and brain injury rehabilitation programs at Parkwood Institute in London, Ontario, and an assistant professor in the Department of Physical Medicine & Rehabilitation at Western University. Her research focuses on the development of prognostic models and predicting outcomes after mild traumatic brain injury and concussion. Most recently, she completed a Master of Science degree in Epidemiology at the Harvard T. H. Chan School of Public Health.



Jo-Anne Aubut BA



Mark Bayley, MD, FRCPC, is Program Medical Director & Psychiatrist-in-Chief at UHN-Toronto Rehabilitation Institute. He is a Professor at the University of Toronto in the Division of Physical Medicine and Rehabilitation, Faculty of Medicine. Mark has held a number of health system leadership roles including Chair of the Canadian Stroke Best Practices committee, the Stroke Evaluation Committee at Corhealth Ontario, and the Vice Chair of the Greater Toronto Area Rehab Network.



Dr. Robert Teasell is Professor of Physical Medicine and Rehabilitation, Schulich School of Medicine and Dentistry, Western University and a Clinical Researcher at Lawson Research Institute in London, Ontario. He is a clinician at Parkwood Institute, St. Joseph's Health Care London.

## Purpose

The Evidence-Based Review of Acquired Brain Injury (ERABI) is a systematic review of the rehabilitation literature of moderate to severe acquired brain injuries (ABI). It is an annually updated, freely accessible online resource that provides level of evidence statements regarding the strength of various rehabilitation interventions based on research studies. The ERABI is a collaboration of researchers in London, Toronto and Ottawa. Our mission is to improve outcomes and efficiencies of the rehabilitation system through research synthesis, as well as from providing the foundational research evidence for guideline development, knowledge translation, and education initiatives to maximize the real-world applications of rehabilitation research evidence.

## Key Concepts

### Acquired Brain Injury

For the purposes of this evidence-based review, we used the definition of ABI employed by the [Toronto Acquired Brain Injury Network](#) (2005). ABI is defined as damage to the brain that occurs after birth and is not related to congenital disorders, developmental disabilities, or processes that progressively damage the brain. ABI is an umbrella term that encompasses traumatic and non-traumatic etiologies.

**TABLE 1 |** Defining Acquired Brain Injury

Included in ABI definition	Excluded from ABI definition
<p><b>Traumatic Causes</b></p> <ul style="list-style-type: none"> <li>• Motor vehicle accidents</li> <li>• Falls</li> <li>• Assaults</li> <li>• Gunshot wounds</li> <li>• Sport Injuries</li> </ul> <p><b>Non-traumatic Causes</b></p> <ul style="list-style-type: none"> <li>• Tumours (benign/meningioma only)</li> <li>• Anoxia</li> <li>• Subarachnoid hemorrhage (non-focal)</li> <li>• Meningitis</li> <li>• Encephalitis/encephalopathy (viral, bacterial, drug, hepatic)</li> <li>• Subdural Hematoma</li> </ul>	<p><b>Vascular and Pathological Incidents</b></p> <ul style="list-style-type: none"> <li>• Intracerebral hemorrhage (focal)</li> <li>• Cerebrovascular accident (i.e., stroke)</li> <li>• Vascular accidents</li> <li>• Malignant/metastatic tumours</li> </ul> <p><b>Congenital and Developmental Problems</b></p> <ul style="list-style-type: none"> <li>• Cerebral Palsy</li> <li>• Autism</li> <li>• Developmental delay</li> <li>• Down’s syndrome</li> <li>• Spina bifida with hydrocephalus</li> </ul> <p><b>Progressive Processes</b></p> <ul style="list-style-type: none"> <li>• Alzheimer’s disease</li> <li>• Pick’s disease</li> <li>• Dementia</li> <li>• Amyotrophic Lateral Sclerosis</li> <li>• Multiple Sclerosis</li> <li>• Parkinson’s disease</li> <li>• Huntington’s disease</li> </ul>

Given that ‘ABI’ can have multiple definitions, studies with an ‘ABI’ population can be equally heterogeneous in terms of the sample composition. Such studies may include any combination of persons with TBI, diffuse cerebrovascular events (i.e., subarachnoid hemorrhage) or diffuse infectious disorders (i.e., encephalitis or meningitis). The vast majority of individuals with ABI have a traumatic etiology; therefore, much of the brain injury literature is specific to TBI. The terms ABI and TBI have been used intentionally throughout ERABI to provide more information about populations where relevant.

### Moderate to Severe Brain Injury

ABI severity is usually classified according to the level of altered consciousness experienced by the patient following injury (Table 2). The use of level of consciousness as a measurement arose because the primary outcome to understand the severity of an injury is the Glasgow Coma Scale. Consciousness levels following ABI can range from transient disorientation to deep coma. Patients are classified as having a mild, moderate or severe ABI according to their level of consciousness at the time of initial assessment. Various measures of altered consciousness are used in practice to determine injury severity. Common measures include the Glasgow Coma Scale (GCS), the duration of loss of consciousness (LOC), and the duration of post-traumatic amnesia (PTA). Another factor used to distinguish moderate and severe brain injury is evidence of intracranial injury on conventional brain imaging techniques which distinguish severity of injury from a mild or concussion related brain injury.



**TABLE 2 |** Defining Severity of Traumatic Brain Injury, adapted from Veterans Affairs Taskforce (2008) and Campbell (2000)

Criteria	Mild	Moderate	Severe	Very Severe
Initial GCS	13-15	9-12	3-8	Not defined
Duration LOC	< 15minutes*	<6 hours	6-48 hours	>48 hours
Duration PTA	< 1hour*	1-24 hours	1-7 days	>7 days
	*This is the upper limit for mild traumatic brain injury; the lower limit is any alteration in mental status (dazed, confused, etc.).			

## Methods

An extensive literature search using multiple databases (CINAHL, PubMed/MEDLINE, Scopus, EMBASE, and PsycINFO) was conducted for articles published in the English language between 1980–December 2018 that evaluate the effectiveness of any intervention/treatment related to ABI. The references from key review articles, meta-analyses, and systematic reviews were reviewed to ensure no articles had been overlooked. For certain modules that lacked research evidence the gray literature, as well as additional databases, were searched in order to ensure the topic was covered as comprehensively as possible.

Specific subject headings related to ABI were used as the search terms for each database. The search was broadened by using each specific database’s subject headings, this allowed for all other terms in the database’s subject heading hierarchy related to ABI to also be included. The consistent search terms used were “head injur\*”, “brain injur\*”, and “traumatic brain injur\*”. Additional keywords were used specific to each module. A medical staff librarian was consulted to ensure the searches were as comprehensive as possible.

Every effort was made to identify all relevant articles that evaluated rehabilitation interventions/treatments, with no restrictions as to the stage of recovery or the outcome assessed. For each module, the individual database searches were pooled, and all duplicate references were removed. Each article title/abstract was then reviewed; titles that appeared to involve ABI and a treatment/intervention were selected. The remaining articles were reviewed in full.

Studies meeting the following criteria were included: (1) published in the English language, (2) at least 50% of the study population included participants with ABI (as defined in Table 1) or the study independently reported on a subset of participants with ABI, (3) at least three participants, (4) ≥50% participants had a moderate to severe brain injury (as defined in Table 2), and (5) involved the evaluation of a treatment/intervention with a measurable outcome. Both prospective and retrospective studies were considered. Articles that did not meet our definition of ABI were excluded.



## Interpretation of the Evidence

The levels of evidence (Table 3) used to summarize the findings are based on the levels of evidence developed by Sackett et al. (2000). The levels proposed by Sackett et al. (2000) have been modified; specifically, the original ten categories have been reduced to five levels. Level 1 evidence pertains to high quality randomized controlled trials (RCTs) (PEDro  $\geq 6$ ) and has been divided into two subcategories, level 1a and level 1b, based on whether there was one, or more than one, RCT supporting the evidence statement.

The evidence statements made in evidence-based reviews are based on the treatment of groups rather than individuals. There are times when the evidence will not apply to a specific case; however, the majority of patients should be managed according to the evidence. Ultimately, the healthcare professional providing care should determine whether an intervention is appropriate and the intensity with which it should be provided, based on their individual patient’s needs. Furthermore, readers are asked to interpret the findings of studies with caution as evidence can be misinterpreted. The most common scenario occurs when results of a trial are generalized to a wider group than the evidence allows. Evidence is a tool, and as such, the interpretation and implementation of it must always be done with the known limitations in mind.

**TABLE 3 | Levels of Evidence**

Level	Research Design	Description
1A	Randomized Controlled Trial (RCT)	More than one RCT with PEDro score $\geq 6$ . Includes within subject comparisons, with randomized conditions and crossover designs
1B	RCT	One RCT with PEDro $\geq 6$
2	RCT	One RCT with PEDro $< 6$
	Prospective Controlled Trial (PCT)	Prospective controlled trial (not randomized)
	Cohort	Prospective longitudinal study using at least two similar groups with one exposed to a particular condition
3	Case Control	A retrospective study comparing conditions including historical controls
4	Pre-Post Trial	A prospective trial with a baseline measure, intervention, and a post-test using a single group of subjects
	Post-test	A prospective intervention study using a post intervention measure only (no pre-test or baseline measurement) with one or more groups
	Case Series	A retrospective study usually collecting variables from a chart review
5	Observational study	Using cross sectional analysis to interpret relations
	Clinical Consensus	Expert opinion without explicit critical appraisal, or based on physiology, biomechanics or “first principles”
	Case Reports	Pre-post or case series involving one subject

## Strength of the Evidence

The methodological quality of each randomized controlled trial (RCT) was assessed using the Physiotherapy Evidence Database (PEDro) rating scale developed by the Centre for Evidence-Based Physiotherapy in Australia (Moseley et al., 2002). The PEDro is an 11-item scale; a point is awarded for ten satisfied criterion yielding a score out of ten. The first criterion relates to external validity, with the remaining ten items relating to the internal validity of the clinical trial. The first criterion, eligibility criteria, is not included in the final score. A higher score is representative of a study with higher methodological quality.

# PRINCIPLES AND MODELS OF CARE

POST ACQUIRED BRAIN INJURY

# Summary of the Evidence

Intervention	Key Point Level of Evidence
<b>Acute Management and Implementation of Care</b>	
Implementation Guidelines	<p>Implementation of TBI protocols based on guidelines from the American Association of Neurologic Surgeons, the Brain Trauma Foundation, or generally accepted best practices may improve mortality and patient outcomes in patients with TBI, however, this may only be the case if protocol compliance is sufficiently high.</p> <p>Implementation of a neurocritical care consult service, introduction of mutual neurocritical care/neurosurgery rounds, introduction of a TBI protocol, and clustering of patients with a neurocritical care diagnosis in the same unit may improve hospital mortality post TBI.</p> <p>Formalized early intervention programs can reduce coma duration, hospital length of stay, improve cognitive levels at discharge and rate of discharges home in patients with TBI.</p> <p><i>There is level 2 evidence that implementation of a protocol based on the American Association of Neurologic Surgeons TBI guidelines may improve mortality compared to patients with TBI prior to guideline implementation.</i></p> <p><i>There is level 2 evidence that implementation of a standard treatment protocol based on generally accepted best practices may decrease mortality and improve discharge Glasgow Outcome Scale scores compared to patients with TBI prior to treatment protocol implementation.</i></p> <p><i>There is level 2 evidence that implementation of a neurocritical care consult service, introduction of mutual neurocritical care/neurosurgery rounds, introduction of a TBI protocol, and clustering of patients with a neurocritical care diagnosis in the same unit may improve hospital mortality compared to prior protocols post TBI.</i></p> <p><i>There is level 2 evidence that a joint commission-certified TBI program may reduce 24 hour and 6 month mortality compared to patients with TBI prior to program implementation.</i></p> <p><i>There is level 2 evidence that implementation of a protocol based on the Brain Trauma Foundation guidelines may reduce mortality in patients with TBI compared to retrospective controls, but only if compliance with the protocol is sufficient.</i></p> <p><i>There is level 3 evidence that a formalized early intervention program may reduce coma duration and length of stay and improve cognitive levels at discharge and percent of discharges to home, compared to extended care facilities in patients with TBI.</i></p>

Intervention	Key Point Level of Evidence
<p>Institutional Comparisons</p>	<p>Greater resource availability and more aggressive care may improve mortality in patients with severe TBI.</p> <p><i>There is level 2 evidence that hospitals that perform computerized axial tomography scans more frequently, use intracranial pressure monitors more often, and admit a higher proportion of patients to the intensive care unit, may have lower severe TBI mortality rates.</i></p> <p><i>There is level 3 evidence that level 1 trauma centers may have lower severe TBI mortality rates than level 2 trauma centers.</i></p>
<p><b>Inpatient Rehabilitation</b></p>	
<p>Functionally based streamed models of inpatient rehabilitation may improve targeted deficits more than all-encompassing traditional inpatient rehabilitation methods in patients with TBI.</p> <p>Compared to a single discipline approach, a coordinated, multidisciplinary approach to inpatient rehabilitation may result in greater functional improvements that are sustained for longer in patients with TBI.</p> <p><i>There is level 3 evidence that functionally-based streamed models of inpatient ABI care, particularly neurophysical and neurocognitive streams, may improve functional independence measure efficiency and reduce disability rating scale scores, respectively, compared to traditional inpatient rehabilitation.</i></p> <p><i>There is level 2 evidence that a coordinated, multidisciplinary inpatient rehabilitation approach may increase the extent of improvement seen in TBI motor and cognitive outcomes as well as lead to longer-term maintenance (up to 24 months) of the treatment effect, compared to single discipline treatment.</i></p>	
<p>Intensity of Inpatient Rehabilitation</p>	<p>Increasing inpatient rehabilitation intensity can reduce hospital length of stay post ABI.</p> <p>Increasing inpatient rehabilitation intensity compared to standard therapy, can improve Glasgow Outcome Scale scores and functional outcomes post ABI in the short term.</p> <p>The efficacy of increased inpatient rehabilitation intensity post ABI can change based on the rehabilitation institution and available resources.</p> <p>Inpatient therapy intensity predicts motor functioning post ABI at discharge.</p> <p>High-level involvement in neurorehabilitation goal setting may result in a greater number of attained goals being maintained at follow-up (two months) in individuals with an ABI.</p>

Intervention	Key Point Level of Evidence
	<p><i>There is level 1b evidence that more intensive inpatient rehabilitation may improve Glasgow Outcome Scale scores at 2 months, but not necessarily at 3 months and beyond, compared to conventional treatment in patients with TBI.</i></p> <p><i>There is level 3 evidence that increasing inpatient rehabilitation intensity may reduce hospital length of stay compared to conventional therapy post ABI.</i></p> <p><i>There is level 2 evidence that inpatient therapy intensity post ABI predicts motor functioning at discharge.</i></p> <p><i>There is level 2 evidence that a high-level of involvement in neurorehabilitation goal setting may result in a greater number of attained goals being maintained at follow-up (two months) compared to neurobehavioral therapy in individuals with an ABI.</i></p>
<p>Timing of Inpatient Rehabilitation</p>	<p>Early inpatient rehabilitation is associated with better outcomes in individuals post ABI.</p> <p>Inpatient rehabilitation in the chronic phase of ABI can still yield meaningful results.</p> <p><i>There is level 2 evidence that inpatient rehabilitation within 35 days post ABI is associated with better outcomes such as shorter comas and hospital length of stay, higher cognitive levels at discharge, better Functional Independence Measure scores, and a greater likelihood of discharge to home, compared to rehabilitation initiated after 35 days post-ABI.</i></p> <p><i>There is level 2 evidence that individuals with an ABI can still benefit from inpatient rehabilitation efforts initiated more than 12 months after sustaining an ABI.</i></p>

**Outpatient Rehabilitation and Community Reintegration**

Earlier outpatient rehabilitation is associated with better outcomes post ABI.

More intensive outpatient rehabilitation is associated with better functional outcomes post ABI. However, this may not be the case if intensity is high enough to interfere with a patient’s ability to perform day to day responsibilities.

Compared to individuals with an ABI who are treated in the community, those treated at an outpatient clinic may be less dependent on support from others, more independent in mobility, display fewer inappropriate social behaviours, and have less difficulty with motor speech.

Outpatient care provided at a residential treatment center may improve motor and cognitive function to a greater extent than when care is provided at a nursing facility or at home in individuals with a TBI.

*There is level 2 evidence that earlier initiation of outpatient and community-based rehabilitation is associated with better functional outcomes compared to residential rehabilitation in individuals with an ABI.*

Intervention	Key Point Level of Evidence
	<p><i>There is level 2 evidence that more intensive outpatient rehabilitation programs are associated with better functional outcomes compared to standard therapy in individuals with an ABI.</i></p> <p><i>There is level 2 evidence that compared to individuals with an ABI who are treated in the community, those treated at an outpatient clinic may be less dependent on support from others, more independent in mobility, display fewer inappropriate social behaviours, and have less difficulty with motor speech.</i></p> <p><i>There is level 1b evidence that a supervised fitness-center based program may be equally as effective as an unsupervised home-based program for improving cardiorespiratory fitness in individuals with an ABI.</i></p> <p><i>There is level 3 evidence that outpatient care provided at a residential treatment center may improve motor and cognitive function to a greater extent than when care is provided at a nursing facility or at home in individuals with a TBI.</i></p>
<p><b>Vocational Rehabilitation</b></p>	<p>Individualized and group vocational rehabilitation programs can improve goal-specific performance and behavioural competency/psychological well-being in individuals post ABI, respectively.</p> <p>Combining specialized vocational rehabilitation services with a community reintegration outpatient group intervention or comprehensive day treatment may not improve community-based employment compared to specialized vocational rehabilitation alone in individuals with an ABI.</p> <p><i>There is level 1b evidence that neither individualized nor general vocational rehabilitation programs may improve performance in goal-specific areas compared to waitlist controls in individuals with ABI.</i></p> <p><i>There is level 2 evidence that combining specialized vocational rehabilitation services with a community reintegration outpatient group intervention or comprehensive day treatment may not improve community-based employment compared to specialized vocational rehabilitation alone in individuals with ABI.</i></p>
<p><b>Complete Care Pathways</b></p>	
<p>Although continuity of care has been shown to be beneficial in optimizing recovery, there is insufficient evidence to draw conclusions regarding the ideal structure of a complete model of ABI care. Further research is required in determining the ideal structure of a complete model of ABI care.</p> <p><i>There is level 2 evidence that individuals post ABI living in both rural and urban settings may have greater functional gains from an integrated network of inpatient, outpatient, and community services compared to standard inpatient rehabilitation.</i></p>	




## Introduction

Acquired brain injury (ABI) presents unique challenges that make rehabilitation difficult to standardize. Further complicating the issue, the development of best-practice principles has been hindered by limited sample sizes and comparison groups in a clinical rehabilitation environment (National Institute of Health, 1998). Additionally, there is a need for more prospective studies that utilize a uniform approach (i.e., standardized assessment and outcome measures) (New Zealand Guidelines Group, 2006). As a result, a consensus on optimal models of care for patients with ABI has been elusive.

In October of 2007, a workshop was held by the National Institute of Neurological Disorders and Stroke to develop a classification system for Traumatic Brain Injury (TBI) designed to direct therapeutic interventions (Saatman et al., 2008). Traditional classification systems have been problematic given the diverse range of care patients require following brain injury. This international group of experts emphasized that this work only begins to scratch the surface in understanding brain injury care. Nevertheless, a model depicting how patients should flow through the health care system following a brain injury has emerged.

Generally after an ABI, patients receive the following in order: pre-hospital care, acute care – which may include an Emergency Department visit, neurosurgical intervention and/or Intensive Care Unit (ICU) management as necessary, inpatient rehabilitation, and then they are discharged to the community with varying levels of support (Khan et al., 2002). Across settings, this pathway may include cognitive and behavioural rehabilitation programs, community living opportunities, rehabilitation services in the home, care management, and prevention initiatives (Zygun et al., 2005). However, despite effective triage programs, best evidence-based protocols, and progress in the management of secondary complications of severe TBI, significant regional differences in practice exist (Zygun et al., 2005). The typical pathway of care is depicted in Figure 3.1; this figure will be replicated throughout the document as the various stages are explained in more detail.



**Figure 1 |** A schematic depiction of the progression of ABI management.

Internationally, rehabilitation care received by patients with brain injury is extremely variable. Care is dictated by local health care policy, local culture, and resource availability. This in turn has made development of internationally applicable systems challenging. In 1965, the World Federation of Neurosurgical Societies formed an “ad hoc” Committee on Head Injuries which was followed by the formation of the Committee of Neuro-traumatology in 1977 (Teasdale et al., 1997). This provided one of

the first published international discussions of brain trauma care. The formation of the International Brain Injury Association in 1993 and the International Association for the Study of Brain Injury in 1998 continued to expand opportunities for the sharing of information (International Brain Injury Association, 2008). In 1995, the Brain Trauma Foundation (BTF) developed the first Guidelines for the Management of Severe TBI (Carney, 2007); these guidelines have since been revised. These guidelines are maintained in conjunction with the American Association of Neurological Surgeons and the Congress of Neurological Surgeons and other stakeholders, such as the European Brain Injury Consortium. Since their inception, countries such as Italy, Mexico, Ireland, and Japan have adapted BTF guidelines to suit local needs (Citerio et al., 2003; Espinosa-Aguilar et al., 2008; Matta & Menon, 1996; Shigemori & Tokutomi, 2002). The World Health Organization has also expanded its focus to assess the need for effective global rehabilitation programs. It has been estimated that over 80% of the world's people with disabilities live in low to middle income countries and only 2% have access to rehabilitation services (Hyder et al., 2007). This is especially relevant when we consider that the highest rates of TBI due to road traffic incidents are in the Latin American and Caribbean regions, with rates in Sub-Saharan Africa not far behind (Hyder et al., 2007). Other countries such as the United States are also seeing an increase in the rates of TBI despite education and prevention (Centers for Disease Control and Prevention, 2018a). From 2001 to 2010, rates of TBIs among males has only increased. In 2001, approximately 600 per every 100 000 males in the US sustained a TBI, and this number has increased to just over 900 per every 100 000 in 2010 (Centers for Disease Control and Prevention, 2018a). Although the rates of TBI in women have also increased over the years, men are still 40% more likely to sustain a TBI (Centers for Disease Control and Prevention, 2018a). In 2014 there were over 2.87 million TBI related emergency department visits, hospitalizations, and deaths in the United States (Centers for Disease Control and Prevention, 2018b).

One of the most comprehensive national brain injury clinical care systems has evolved in the United States (US). In 1978 the National Institute on Disability and Health Research (now the National Institute on Disability and Rehabilitation Research) provided funding to New York University's Rusk Center and the Santa Clara Valley Medical Center (San Jose, California) to develop a model of dedicated, interdisciplinary, acute inpatient rehabilitation, coupled with post-acute rehabilitation interventions and cognitive and behavioural approaches (Cope et al., 2005). By August 2004, ABI care in the US included 123 accredited hospitals, 9 skilled nursing facilities (acute inpatient rehabilitation), 153 outpatient programs, 51 home and community programs, 212 long-term residential programs, 231 residential programs and 86 vocational programs (Cope et al., 2005). While there is no one body which oversees brain injury rehabilitation specifically, several organizations have been developed in an attempt to improve the cohesion of the system. Some of the more influential organizations include the Brain Injury Association of America which was established in 1980 and currently works with 40 state run brain injury affiliates (Brain Injury Association of America, 2015) to provide community services to individuals with brain injuries. The National Association of State Head Injury Administrators developed in 1990 as a forum to provide information to State governments and policy makers regarding brain injury (National Association of State Head Injury Administrators, 2008 ), while the Center for Disease Control collects epidemiological information and sponsors research through the Public Health Injury Surveillance and Prevention Program (Centers for Disease Control, 2008). Finally, the Traumatic Brain Injury Model Systems of Care was developed in 1997 as a prospective, longitudinal multi-center study to assess

rehabilitation of patients through a coordinated system of acute care and inpatient rehabilitation with a 15 year follow-up (National Data and Statistical Center, 2008). Although these four organizations and others like them work together to provide guidance regarding brain injury care, ultimately decisions are still left to individual institutions and their clinicians, resulting in regional differences in care.

In Canada, brain injury rehabilitation has steadily developed in a similar way to the American system. During the 1980's and 90's, brain injury rehabilitation evolved as a specialization of rehabilitation medicine. Rehabilitation hospitals work within provincial health care systems and, as a result, some provinces, particularly the more scarcely populated ones, have more limited ABI rehabilitation resources. Moreover, within provinces there is often a disparity in services between larger urban centers and smaller rural areas. While access to care is universally available, private services can also be utilized by those patients with access to private funding (Cullen, 2007). Reid-Arndt et al. (2010) explain that whilst patients have greatly benefitted under the Rehabilitation Act after the passing of the TBI Act in 1996, community-based interventions, employment services, and independent living programs continue to require additional funding and support.

In 2003, the Brain Injury Association of Canada was established to provide a national forum for sharing brain injury information. Currently, only the territories lack a territorial level brain injury association (Brain Injury Association of Canada, 2015). In an attempt to standardize care, Accreditation Canada, a not-for-profit organization, assesses health care institutions in Canada for quality of care and specifically includes brain injury services (Accreditation Canada, 2008). The Canadian Institute for Health Information was established by national, provincial, and territorial governments to collect and disseminate health information including information regarding rehabilitation facilities. Rehabilitation information is drawn from the National Rehabilitation Reporting System with 99 facilities across nine provinces submitting data (Canadian Institute for Health Information, 2014). A separate database has also been established at the Toronto Rehabilitation Institute, which is modeled after the American Model systems. The Canadian database was expanded in 2002 to uniquely include individuals with non-traumatic brain injuries as well, which differs from the American system (Cullen, 2007).

Europe presents some unique cultural and political challenges in brain injury. The European Brain Injury Society was formed in 1989 and now has 181 institutional members from all nations in the European Union, as well as Switzerland (European Brain Injury Society, 2015). The European Brain Injury Consortium was formed in 1994, which *“reflected the realization that numbers of patients required in the design of definitive Phase III studies of severe head injury demanded European-wide recruitment”* (p.798) (Teasdale et al., 1997). While nations were encouraged to continue to develop their own strategies, value was placed on international collaboration. In 1997 the European Brain Injury Consortium developed guidelines for management of severe head injury in adults to attempt to provide some clarity and standardization in brain injury care (Maas et al., 1997). With similar collaborative goals, the European Brain Council was formed in 2002 in Brussels to attempt to coordinate research in the area of brain disease, including brain injury (Olesen & Freund, 2006). Despite these attempts at standardization, national models of ABI care are still dictated by regional health care policies.

March et al. (2013) reviewed the National Mental Health Plan and reported that ABI was not referred to or explicitly acknowledged and that ABI service providers were not incorporated into future target partnerships. Given how common mental health concerns are within the ABI population, March et al. (2013) argue that this absence of acknowledgement hinders dual-diagnosis care and that a collaborative approach with stakeholders will help provide leadership in addressing this issue.

With a global perspective in mind, this module presents a broad analysis of the over-riding systems of care in ABI. Papers were considered for analysis if they focused on a generalized system of care. Since the aim of this module is to compare different models of rehabilitation and not assess the effectiveness of individual rehabilitation strategies by themselves, only papers that compare at least two distinct rehabilitation groups were included. These could include separate hospitals, separate treatment groups within one center, or comparisons between patients in the same center before and after systemic changes.

## Acute Management and Implementation of Care



The most severe consequences of an ABI are often not due to the initial trauma itself. Secondary brain injury can result from edema, ischemia, elevated intracranial pressure, and inadequate cerebral perfusion pressure, as well as calcium imbalances, excitatory amino acid release, and free radical production; all of which can lead to cell death (Zasler et al., 2007). For this reason, the speed and intensity with which patients are cared for is of the utmost importance. Assessments of how to acutely treat patients with ABI generally fall into one of four categories: pre-hospital care, hospital facility type, adherence to acute care guidelines, and discharge destination. Each of these areas presents a unique challenge. Module 14 of this evidence-based review reports the current evidence for acute treatment of ABI. Here we have attempted to highlight concerns and elucidate attempts being made to improve the current system of care with reference to the application and efficacy of treatment protocols and guidelines, and by comparing models of care across different institutions.

Pre-hospital care can be the difference between life and death. The time from injury to intervention is perhaps the most obvious component of pre-hospital care but debate has also arisen regarding the types of treatments that are suitable prior to hospital arrival. In 2000, the BTF released guidelines for pre-hospital management of patients with ABI. An Emergency Medical Service task force developed a

consensus based algorithm (Gabriel et al., 2002). The guidelines were then updated in 2007 (Badjatia et al., 2007). The French guidelines pertaining to the first 24 hours of management following severe TBI were published (Geeraerts et al., 2018). Nevertheless, the degree of variability in the way acute care is delivered in the US is still fairly unknown (Bulger et al., 2007). This has also been shown to be true of other countries that have begun to examine protocols for out-of-hospital care (Baethmann et al., 1999; Harrington et al., 2005). Research has been conducted regarding the efficiency of transfer and access to trauma centers in general (Bulger et al., 2007) but little to no research has been performed specifically for brain injury.

The final stage of acute care involves the transition to post-acute care. Once patients are medically stable, they are transferred to one of three places: home, long term care, or a rehabilitation unit. Rehabilitation units for patients with ABI can consist of hospital-based inpatient rehabilitation centers or specialized rehabilitation units that often focus on behavioural issues. One study has shown that approximately 85% of patients are transferred to inpatient rehabilitation within 1 year of ABI (Godbolt et al., 2015). How and by whom this decision is made may greatly affect the type of care that is received by patients. Several factors, such as availability of rehabilitation inpatient beds, the patient's support needs, and the patient's financial situation may play a role in this decision. In the US, Medicaid patients were 68% and Health Maintenance Organization patients were 23% more likely to be discharged to a skilled nursing facility than those on a fee-for-service plan (Chan et al., 2001). In Canada, patients injured in a motor vehicle accident were 1.6 times more likely to be discharged home with support services than those who were injured in a fall (Kim et al., 2006). The standardization of care across multiple institutions helps to ensure that each individual is receiving the best available level of care, and that it is being applied in the most effective manner.

## Implementation of Guidelines

As mentioned earlier, guidelines have been established by organizations such as the BTF and the European Brain Injury Consortium to standardize treatment and to aid in the dissemination of information. Audits of guideline implementation can help to ensure that a proper level of care is provided in all types of medical centers.

Juelsgaard et al. (2018) assessed the degree to which the Danish Helicopter Emergency Medical Service critical care teams adhered to the prehospital guidelines for the treatment of patients with acute non-traumatic intracranial pathology or isolated TBI. They discovered complete adherence to recommendations (endotracheal intubation, SpO<sub>2</sub> > 90%, and SBP > 90mmHg) in 69% of non-TBI cases and 74% of isolated TBI cases.

Hesdorffer and Ghajar (2007) demonstrated that there was an increasing trend in guideline adherence with nurses and Intensive Care Unit trauma coordinators. This study showed that guideline adherence improved from 17% to 44.7% from 2000-2006. In the US alone, it is estimated that a modest improvement to 50% adherence of BTF guidelines from 33% would result in 989 lives saved annually (Faul et al., 2007). A study by Griesdale et al. (2015) revealed that adherence to BTF guidelines regarding

cerebral perfusion pressure was low with recommended pressure ranges occurring only 31.6% of the time. In addition, Talving et al. (2013) also reported that compliance with BTF guidelines for monitoring intracranial pressure (ICP) was at 46.8%, resulting in significantly higher mortality and brain herniation among patients not receiving ICP monitoring (Peep Talving et al., 2013). Shafi et al. (2014) revealed a mean compliance rate of 73% to the BTF guidelines, with only 3 of 11 level-1 trauma centres achieving rates of over 80%. Compliance varied widely among centers reflecting that patients with severe TBI receive inconsistent care. A systematic review by Khormi et al. (2018), reported the median percentage of adherence to the BTF guidelines as 60.7% with no significant difference in adherence level noted for studies performed in North America when compared with those done in other countries. Unfortunately, this statistic has not improved since a 60% adherence level was observed in 2002. Interestingly, Khormi et al. (2018) also found that the level of adherence was proportional to the strength of evidence, with level I recommendations associated with more optimal adherence. Furthermore, organizational factors such as treatment in a Level I trauma center and living in a country with higher economic status positively influenced adherence, whereas lack of health insurance negatively influenced it.

An online survey administered by Lamontagne et al. (2018) revealed that only 47% of respondents knew of at least one CPG to support the rehabilitation of individuals following moderate-severe TBI, however, only 34% of the documents named by respondents met criteria as a CPG. In addition, 53% of respondents said they did not use a CPG to guide their practice. Respondents identified lack of time and/or staff as well as a heavy caseload as the top reasons why they did not feel well-equipped to use new CPGs. Respondents indicated a preference for specific, as opposed to general, recommendations as well as a ranked list of recommendations based on the associated level of evidence. Furthermore, they requested that recommendations be accessible electronically and online rather than through more traditional paper format.

Lamontagne et al. (2018) asked providers in Quebec and Ontario to report their perceived level of implementation for a subset of key recommendations contained within the INESSS-ONF CPG for the rehabilitation of adults with moderate-severe TBI. Overall, the proportion of recommendations perceived as “partially” or “not” implemented was low and varied from 8-34% across guideline subsections. In general, these recommendations were deemed as high or medium priority and felt feasible to implement by respondents. Recommendations in the “Components of the Optimal TBI Rehabilitation System” section were less implemented than those in the “Assessment and Rehabilitation of Brain Injury Sequelae” section. This finding was thought to reflect the difficulty that exists in making system-level changes. It is important to note that this study presents perceived levels of implementation, rather than actual observed levels.

Adherence to guidelines is a continuous process and has a direct impact on patient care. This section summarizes the available evidence evaluating the effect of guideline/protocol implementation on patient outcomes in ABI.



**TABLE 4 |** Guideline Implementation for Acute Management Post ABI

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Sorinola et al.</a> (2018) Hungary PCT N= 7230</p>	<p><b>Population:</b> <i>severe TBI Group (N= 7230):</i> Mean Age= 60.89yr; Gender: Male=72%, Female=28%. <b>Intervention:</b> Health records of TBI individuals were examined of at hospitals in Hungary. Authors investigated the impact of a TBI guideline introduction. Authors classified the 8 institutions that provided over 50% of the care as ‘centers’ and the remaining 49 as ‘secondary institutions.’ <b>Outcomes:</b> Case Fatality Ratios (CFR).</p>	<ol style="list-style-type: none"> <li>1. Pre-guideline introduction saw CFRs of 23.4%, 37.7%, and 47% at 1wk, 1mo, and 6mos, respectively at the centers.</li> <li>2. Post-guideline introduction saw CFRs of 22.1%, 39.1%, and 50% at 1wk, 1mo, and 6mo, respectively at the centers.</li> <li>3. Pre-guideline introduction saw CFRs of 21.5%, 39.1%, and 50% at 1wk, 1mo, and 6mo, respectively at the secondary institutions.</li> <li>4. Post-guideline introduction saw CFRs of 21.9%, 37%, and 48.9% one week, one month, and 6 months, respectively at the secondary institutions.</li> <li>5. The center and secondary institution CFRs showed no significant change when comparing the pre and post guideline periods.</li> </ol>
<p><a href="#">Tarapore et al.</a> (2016) The Netherlands Cohort N=832</p>	<p><b>Population:</b> TBI; Gender: Male=654, Female=178; Mean Age=38yr; GCS: mild (n=178), moderate (n=118), severe (n=466). <b>Intervention:</b> A joint-commission-certified TBI program was implemented at San Francisco General Hospital and patient outcomes were compared to historic controls. <b>Outcome Measure:</b> Early TBI deaths (&lt;24 hr), mortality at 6 mo.</p>	<ol style="list-style-type: none"> <li>1. The percentage of early TBI deaths (&lt;24hr) were 59% lower in the Joint Commission-certified TBI program cohort as compared to the historical control cohort.</li> <li>2. The percentage of observed deaths 6mo after the Joint Commission-certified TBI program was instituted was 22% lower as compared to the historical cohort.</li> </ol>
<p><a href="#">Kesinger et al.</a> (2014) USA Case Control N=108</p>	<p><b>Population:</b> TBI; <i>Pre-Standard Trauma Protocols (STP; n=68):</i> Mean Age=37.1yr; Gender: Male=63, Female=5. <i>Post-STP (n=40):</i> Mean Age=38.6yr; Gender: Male=31, Female=9. <b>Intervention:</b> Chart reviews were conducted comparing hospital records pre and post implementation of STPs. These protocols were based on best practices and damage control resuscitation (e.g., small volume resuscitation, requiring a physician’s presence in intra-hospital transportation of severely injured patients, etc.). <b>Outcome Measure:</b> Glasgow Coma Scale (GCS), Medical Interventions, Mortality Rates.</p>	<ol style="list-style-type: none"> <li>1. In the emergency department, after STPs were implemented, there was an increase in resuscitation with 7.5% hypertonic saline (p=0.014), use of catheters (p=0.015), administration of tetanus vaccinations (p=0.034), and earlier use of blood transfusions (p=0.008).</li> <li>2. Post STP, hospital mortality decreased from 38% to 18% (p=0.024) and GCS scores improved from a median of 10 to a median of 14 (p=0.034).</li> </ol>



Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Kramer &amp; Zygun</a> (2013) Canada Cohort N=4,097	<p><b>Population:</b> TBI=1604, Anoxic BI=552, Subarachnoid Hemorrhage=449, Intracerebral Hemorrhage= 398, Stroke=444, Central Nervous System Infection=242, Status Epilepticus=605; Gender: Male=2581, Female=1516.</p> <p><b>Intervention:</b> Patient data was extracted from an Intensive Care Unit (ICU) database over four time periods based on when new protocols were developed and introduced. New protocols included: Neurocritical Care Consult Service (September 2003), Temperature Regulation Protocol (September 2004), Mutual Neurocritical Care/Neurosurgery Rounds (July 2005), TBI Protocol (August 2008), and Clustering of Neurocritical Care Patients (June 2010).</p> <p><b>Outcome Measure:</b> Hospital Mortality, Discharge Home Without Support.</p>	<ol style="list-style-type: none"> <li>Hospital mortality improved significantly after implementation of a Neurocritical Care Consult Service (p=0.03; Odds ratio=0.81), Mutual Neurocritical Care/Neurosurgery Rounds (p=0.008; Odds ratio=0.80), TBI Protocol (p=0.04; Odds ratio=0.84), and Clustering of Neurocritical Care Patients (p=0.02; Odds ratio=0.76) and improved non-significantly after implementation of the Temperature Regulation Protocol (p=0.07; Odds ratio=0.85).</li> <li>Discharge without home support improved significantly after implementation of a Neurocritical Care Consult Service (p=0.04; Odds ratio=1.27), Mutual Neurocritical Care/Neurosurgery Rounds (p=0.0002; Odds ratio=1.39), Clustering of Neurocritical Care Patients (p=0.01; Odds ratio=1.31), and implementation of the Temperature Regulation Protocol (p=0.0009; Odds ratio=1.38), and improved non-significantly after TBI Protocol implementation (p=0.06; Odds ratio=1.17).</li> </ol>
<a href="#">Myburgh et al.</a> (2008) Australia/NZ Cohort N=635	<p><b>Population:</b> TBI; Mean Age=41.6yr; Gender: Male=471, Female=164; Severity: Mild=159, Moderate=114, Severe=362.</p> <p><b>Intervention:</b> Data was obtained prospectively for patients cared for after the publication of new Brain Trauma Foundation guidelines and compared to retrospective control data (pre-guidelines). Follow-up telephone interviews were conducted at 6mo and 12mo post-injury.</p> <p><b>Outcome Measure:</b> Glasgow Outcome Scale Extended (GOSE), Mortality.</p>	<ol style="list-style-type: none"> <li>Favourable outcomes on the GOSE were found in 58.8% of all patients, and 48.5% of patients with severe TBI.</li> <li>Mortality was reported in 26.9% of all patients and 35.1% for patients with a severe TBI.</li> <li>Although concordance with guideline management was generally seen; mortality and favorable neurological outcomes were similar to previous studies before the advent of evidence-based guidelines.</li> </ol>
<a href="#">Fakhry et al.</a> (2004) USA Case Control N=830	<p><b>Population:</b> TBI; <i>Group 1 (n=219):</i> Mean Age=33.8yr; Gender: Male=161, Female=58; Mean GCS=4. <i>Group 2 (n=188):</i> Mean Age=33.9yr; Gender: Male=133, Female=55; Mean GCS=3.5. <i>Group 3 (n=423):</i> Mean Age=35.6yr; Gender: Male=327, Female=96; Mean GCS=3.5.</p> <p><b>Intervention:</b> Patients were retrospectively divided into 3 groups: Group 1, 1991-1994 (pre-guidelines); Group 2, 1995-1996 (post-guidelines low compliance); and Group 3, 1997-2000 (post-guidelines high compliance). Data was extracted from hospital trauma registries and from chart reviews.</p> <p><b>Outcome Measure:</b> Mortality, length of stay, total charges, Rancho Los Amigos Levels of Cognitive Functioning Scale (RLAS), and Glasgow Outcome Scale (GOS).</p>	<ol style="list-style-type: none"> <li>Significant differences were seen between groups, with group 2 achieving higher GOS scores at discharge (p&lt;0.001), a decrease in length of stay in hospital (p=0.001) and a decrease in total charges per patient (p=0.002).</li> <li>A significant drop in mortality was seen in the Group 3 compared to Group 1 (p=0.047).</li> <li>Appropriate responses on the RLAS significantly improved over time from 43.9% in Group 1, 44% in Group 2 and 56.6% in Group 3 (p=0.004).</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Palmer et al.</a> (2001) USA Case Control N=93	<p><b>Population:</b> TBI; <i>Group 1 (n=37):</i> Mean Age=41.35yr; Mean GCS=6.43. <i>Group 2 (n=56):</i> Mean Age=38.10yr; Mean GCS=6.88.</p> <p><b>Intervention:</b> Data was collected from the medical records of patients with TBI treated pre- (group 1) and post- (group 2) guideline implementation.</p> <p><b>Outcome Measure:</b> Glasgow Outcome Score (GOS) and cost.</p>	<ol style="list-style-type: none"> <li>1. Patients in group 2 were more likely to demonstrate significant gains on GOS (p&lt;0.005) compared to group 1.</li> <li>2. Guideline implementation resulted in a 9.13 times higher odds ratio of good outcome relative to poor outcome or death pre-implementation.</li> <li>3. Hospital charges increased by \$97,000 per patient.</li> </ol>
<a href="#">Mackay et al.</a> (1992) USA Case Control N=38	<p><b>Population:</b> TBI; <i>Group 1 (n=17):</i> Mean Age=29.1yr; Gender: Male=12, Female=5; Mean GCS=5.18. <i>Group 2 (n=21):</i> Mean Age=30yr; Gender: Male=19, Female=2; Mean GCS=5.80.</p> <p><b>Treatment:</b> Chart reviews were completed on patients treated in hospital. Patients in Group 1 received a formalized early intervention program and were compared with patients in Group 2 who were treated at hospitals without a formalized early intervention program.</p> <p><b>Outcome Measure:</b> Length of coma, length of stay, Rancho Los Amigos Scale of Cognitive Functioning and Injury Severity Score.</p>	<ol style="list-style-type: none"> <li>1. Patients in Group 1 experienced shorter comas (p=0.033), lengths of stay (p=0.026) and had a greater likelihood of being discharged to home (94% versus 57%).</li> <li>2. Patients in Group 1 demonstrated significantly greater functional improvements at discharge in cognitive/language levels (p=0.018), motor/physical abilities (p=0.032) and perceptual/sensory skills (p=0.025) compared with Group 2.</li> </ol>

## Discussion

A number of studies have investigated the effect of the implementation of protocols incorporating evidence-based guidelines. Kesinger et al. (2014) implemented a Standard Trauma Protocol (STP) at a level 1 trauma center in Colombia and compared outcomes pre- and post-guideline implementation. After STP implementation, hospital mortality decreased by 20%, and median GCS scores at discharge improved from 10 to 14. In another study, Palmer et al. (2001) measured the impact of a protocol based on the American Association of Neurologic Surgeons traumatic brain injury guidelines in a community hospital. Patients had 9.13 times higher odds ratio of good outcomes relative to the odds of poor outcomes or death in the post-protocol group versus the pre-protocol group. Hospital charges did increase by \$97,000 per patient, however, the authors claim this was justified by the significant improvements in patient outcomes. In southern Alberta, Kramer & Zgun (2013) conducted a cohort study of 4,097 participants over a more than 11-year period to compare patient outcomes before and after four new protocols were introduced. Hospital mortality improved after implementation of a neurocritical care consult service in 2003, introduction of mutual neurocritical care/neurosurgery rounds in 2005, introduction of a TBI protocol in 2008, and clustering of patients with a neurocritical care diagnosis within a larger unit in 2010. Trends also show improvement in hospital mortality after implementation of the temperature regulation protocol in 2004.

While the implementation of protocols has been shown to improve patient outcomes in a number of studies, adherence to protocols is also an important factor to consider. In Hungary, Sorinola et al. (2018) demonstrated how the simple existence of guidelines does not in itself result in improved outcomes for

patients with severe traumatic brain injury. This study highlights the importance of introducing audit processes and measures to enforce guideline compliance. A study by Saherwala et al. (2018) found a dedicated and structured educational program – the Adam Williams Initiative – was associated with high rates of adherence to BTF recommendations. In addition, more specific guidelines were more likely to be adhered to.

Three studies consider both implementation and concordance with protocols in TBI care. In a cohort study, Myburgh et al. (2008) prospectively recorded data for patients cared for after the publication of the 1996 BTF guidelines and compared them to patients treated before. Despite guideline publication, mortality and favorable neurological outcomes remained similar pre- and post-publication. In another study, while concordance with the guidelines was evident for measures such as thromboprophylaxis and head elevation, ICP monitoring was used in only 44.5% of patients with severe TBI, indicating that guideline adherence was lacking. Tarapore et al. (2016) studied the effect of the implementation of a joint-commission certified TBI program at the San Francisco General Hospital (an urban level 1 trauma center). After implementation of the program in 2011, patient data was tracked from 2011-2013 and expected versus observed mortality was compared to a patient database from 1987-1996 which acted as a historical control. The 2011 TBI program reduced the percentage of early mortality by 59%, and 6-month mortality by 22%, compared with the historical control. However, the authors suggest that adopting evidence-based guidelines was not sufficient; they also tracked compliance to ensure that the guidelines were implemented properly. Fakhry et al. (2004) compared patient outcomes before and after implementation of a protocol in 1995 following the BTF guidelines. After implementation, initial analysis indicated compliance with the protocol was only about 50%, however, by 1997, compliance had improved to 88%. Thus, the authors evaluated length of stay and total charges per patient during 3 time periods: 1991-1994 (pre-protocol), 1995-1996 (post-protocol, low compliance), and 1997-2000 (post-protocol, high compliance). The authors found that length of stay and total charges per patient dropped in each of the time periods measured, and mortality was decreased between the 1991-94 and 1997-2000 cohorts. These results indicate that both the implementation of guidelines and the ability to comply with them can make significant differences when it comes to patient outcomes.

TBI care in the acute setting is vital due to the risk of secondary brain injury resulting from increased intracranial pressure or insufficient cerebral perfusion pressure. Since healthcare practitioners have many physiological factors and treatment options to consider when devising a treatment plan, all while being under the pressure of time, an evidence-based, standardized treatment protocol may be particularly warranted in the acute setting. One study examined the efficacy of a formalized TBI early intervention program in comparison to hospitals that lack formalized TBI programming (Mackay et al., 1992). The authors found that the implementation of a formalized early intervention program resulted in significantly lower length of comas, length of stays, significantly higher mean cognitive levels at discharge, and a significantly higher percentage of discharges to home versus extended care facilities.

## Conclusions

*There is level 2 evidence that implementation of a protocol based on the American Association of Neurologic Surgeons TBI guidelines may improve mortality compared to patients with TBI prior to guideline implementation.*

*There is level 2 evidence that implementation of a standard treatment protocol based on generally accepted best practices may decrease mortality and improve discharge Glasgow Outcome Scale scores compared to patients with TBI prior to treatment protocol implementation.*

*There is level 2 evidence that implementation of a neurocritical care consult service, introduction of mutual neurocritical care/neurosurgery rounds, introduction of a TBI protocol, and clustering of patients with a neurocritical care diagnosis in the same unit may improve hospital mortality compared to prior protocols post TBI.*

*There is level 2 evidence that a joint commission-certified TBI program may reduce 24 hour and 6-month mortality compared to patients with TBI prior to program implementation.*

*There is level 2 evidence that implementation of a protocol based on the Brain Trauma Foundation guidelines may reduce mortality in patients with TBI compared to retrospective controls, but only if compliance with the protocol is sufficient.*



### KEY POINTS

- Implementation of TBI protocols based on guidelines from the American Association of Neurologic Surgeons, the Brain Trauma Foundation, or generally accepted best practices may improve mortality and patient outcomes in patients with TBI, however, this may only be the case if protocol compliance is sufficiently high.
- Implementation of a neurocritical care consult service, introduction of mutual neurocritical care/neurosurgery rounds, introduction of a TBI protocol, and clustering of patients with a neurocritical care diagnosis in the same unit may improve hospital mortality post TBI.
- Formalized early intervention programs can reduce coma duration, hospital length of stay, improve cognitive levels at discharge and rate of discharges home in patients with TBI.

## Institutional Comparisons

Facility type is also of prime interest relative to the specific needs of the patient. Trauma care facilities have proven to be superior to general care facilities for emergency medical care. MacKenzie et al. (2007) noted patients with head injuries and an Abbreviated Injury Scale score  $\geq 3$  showed a 90% survival rate at 12-month follow up in trauma centers, compared to 64.3% in non-trauma centers. The availability of trauma centers tends to be dictated by local needs and resources. In the absence of such a facility, local

centers must be able to handle individuals with ABI effectively and transport them when necessary to a properly equipped center. This section provides an overview of studies assessing institutional differences in ABI care.

**TABLE 5 | Institutional Comparisons for Acute Management Post ABI**

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Harris et al.</a> (2008) USA Cohort N=1,607	<p><b>Population:</b> TBI; <i>Hospital 1 (n=691)</i>: Mean Age=38.3yr; Gender: Male=511, Female=180; Severity: Mild=414, Moderate=57, Severe=161. <i>Hospital 2 (n=782)</i>: Mean Age=34.8yr; Gender: Male=633, Female=149; Severity: Mild=324, Moderate=77, Severe=91. <i>Hospital 3 (n=134)</i>: Mean Age=34.6yr; Gender: Male=109, Female=25; Severity: Mild=82, Moderate=19, Severe=17.</p> <p><b>Intervention:</b> Data from three hospitals that provide neurosurgical care was extracted from a National Trauma Registry. Hospital 1 was in the USA and Hospitals 2 and 3 in Jamaica.</p> <p><b>Outcome Measure:</b> Medical intervention use, Mortality rates, Glasgow Outcome Scale (GOS), and Functional Independence Measure (FIM).</p>	<ol style="list-style-type: none"> <li>1. Patients cared for in hospital 1 had more severe head injuries, received more CT scans (<math>p&lt;0.0001</math>), and were more likely to be admitted to the ICU than those in hospitals 2 and 3 (<math>p&lt;0.0001</math>).</li> <li>2. Patients in hospital 1 were more likely to receive intracranial pressure monitoring.</li> <li>3. There were no statistically different differences in mortality rates between the three sites, except severe patients cared for in the USA had a decreased risk of mortality (OR 0.47, <math>p=0.04</math>).</li> <li>4. Patients cared for in the USA had lower mean GOS scores (<math>p&lt;0.0001</math>) and lower FIM scores for self-feed (<math>p=0.0003</math>), locomotion (<math>p=0.04</math>), and verbal (<math>p&lt;0.0001</math>).</li> </ol>
<a href="#">DuBose et al.</a> (2008) USA Case Control N=16,035	<p><b>Population:</b> TBI; Mean Age=40.7yr; Gender: Male=11,169, Female=4866.</p> <p><b>Intervention:</b> Data on patients managed in level I and level II trauma centers was extracted from the National Trauma Data Bank and compared.</p> <p><b>Outcome Measure:</b> Injury Severity Scale (ISS), Mortality, Medical complications, and clinical procedures.</p>	<ol style="list-style-type: none"> <li>1. After adjustments for patient differences, those managed in a level 2 trauma center had increased mortality, more complications and greater likelihood of progression of neurologic insult (all <math>p&lt;0.001</math>).</li> <li>2. <math>ISS\geq 20</math>, <math>Age\geq 55</math>, <math>GCS\leq 8</math>, admission to Level 2 trauma centers, penetrating mechanism and hypotension on admission were all significant risk factors for mortality (all <math>p&lt;0.001</math>).</li> </ol>

## Discussion

Strategies for the management of ABI care vary widely depending on accessibility to resources and the patient population that the healthcare institution serves. Comparing outcomes between institutions with different management strategies may provide much needed information as to what practices are effective at specific levels of resource availability. Two studies were found that compare outcomes for patients with TBI across healthcare institutions that have markedly different management strategies.

Harris et al. (2008) followed TBI care in one US hospital and two Jamaican hospitals to evaluate TBI management strategies in developed versus developing countries. The hospital in the US had more advanced technology and provided more aggressive neurological support. Specifically, the US hospital performed CT scans more frequently, used intracranial pressure monitors more often, and admitted a higher proportion of patients to the ICU. Interestingly, despite the greater availability of resources and aggressiveness of treatment at the US site, the overall mortality rates did not differ between hospitals.

Notably, however, in the subset of patients with severe head injury, mortality was significantly reduced at the US site. Patients with more severe injuries may benefit from more aggressive treatment strategies.

DuBose et al. (2008) aimed to compare severe TBI patient outcomes between trauma centers with different designations. The American College of Surgeons (ACS) (1999) designates trauma centers into 1 of 3 levels according to available resources, education, and research. Previous studies of severe trauma patients have shown that level 1 trauma centers have better mortality and morbidity rates than their level 2 counterparts (Demetriades et al., 2005). In this study, a total of 16,037 patients with severe TBI were split into two groups based on the center in which they were treated (level 1 or level 2). Patients treated at the level 1 trauma center had lower mortality rates, lower rates of complications, and less progression of neurologic insult.

### Conclusions

*There is level 2 evidence that hospitals that perform computerized axial tomography scans more frequently, use intracranial pressure monitors more often, and admit a higher proportion of patients to the intensive care unit, may have lower severe TBI mortality rates.*


*There is level 3 evidence that level 1 trauma centers may have lower severe TBI mortality rates than level 2 trauma centers.*



#### KEY POINTS

- Greater resource availability and more aggressive care may improve mortality in patients with severe TBI.

## Inpatient Rehabilitation



The rehabilitation of patients with ABI involves a comprehensive effort by an interdisciplinary team including physicians, nurses, physical therapists, speech-language pathologists, occupational therapists, and social workers. Considering the incidence, consequences, and costs of ABI, it is important to understand the relative efficacy of rehabilitation methods.

While many patients with ABI are discharged directly home or to a long-term care facility, others are discharged to a dedicated inpatient rehabilitation service. These services vary from institution to



institution but generally include some type of intensive therapy program for physical, social, behavioural, and cognitive difficulties. Deciding who should receive inpatient rehabilitation remains a major challenge. Patient referral decisions are inherently complex and need to be understood as a dynamic phenomenon shaped by characteristics of the individual. However, they also rely on the interactions and interpretations of health professionals who operate within unique organizational and broader health care contexts (Foster & Tilse, 2003). These decisions are also influenced by social and funding issues. For example, in the US, patients insured by Medicaid or a Healthcare Maintenance Organization were more likely to go to a skilled nursing facility, rather than inpatient rehabilitation, relative to people with commercial fee-for-service plans (Chan et al., 2001). In Canada, patients aged 36 to 45 years old with more co-morbid conditions are more likely to receive rehabilitation than those older than 65, rural dwellers, non-English speaking people and individuals with mental health, alcohol and/or drug problems (Colantonio et al., 2004). The diversity of patient needs has also led to the formation of differing systems of rehabilitation. In Calgary, for instance, the Halvar Johnson Centre has established a slow stream rehabilitation program for individuals with TBI who may require slightly extended care.

Due to the unique challenges posed by ABI, the structure of inpatient rehabilitation is extremely diverse. Patients are generally rehabilitated in one of two centers: a general rehabilitation unit or a coordinated multidisciplinary neurorehabilitation unit. Some argue that an effective rehabilitation service requires a multidisciplinary team, which includes nursing care, physician monitoring, psychologist and social work intervention, physiotherapists, occupational therapists, and speech language pathologists, among other things (Cifu et al., 2003). In reality, differences in care occur due to the limited availability of neuro-rehabilitative beds and facilities. Limited resources mandate that decisions be made regarding which patients will benefit most from inpatient rehabilitation compared to community-based programs.

Debate also exists about appropriate targets of rehabilitative care. Traditional rehabilitation models in other disciplines such as stroke, spinal cord, and polio have focused on orthopedic and neuromotor impairments (Cope et al., 2005). Brain Injury rehabilitation initially followed a similar path, however, an increased focus was placed on cognitive and behavioural remediation (Mazaux & Richer, 1998), as well as coma stimulation (Cope et al., 2005). Patients in need of skill application training are often discharged to community based services while inpatient rehabilitation focused more on intensive, short term physical or cognitive rehabilitation (Evans, 1997). Furthermore, some inpatient facilities have recognized the need to divide patients into different streams during rehabilitation. At the Toronto Rehabilitation Institute, for example, patients have been streamed into a Neurocognitive group and a Neurophysical group since 2002 (Cullen, 2007).

Inpatient rehabilitation typically begins when a patient is medically stable enough to be transferred out of acute care and into a dedicated rehabilitation unit for a defined period of interdisciplinary rehabilitation. There is a great deal of variability in the length, type, and intensity of services provided in programs throughout the world. As such, we delineate the evidence supporting the various aspects of treatment for inpatient care delivery (Table 6).

**TABLE 6 |** Models of Inpatient Rehabilitation for ABI



Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Cullen et al.</a> (2013) Canada Case Control N<sub>initial</sub>=138, N<sub>final</sub>=130</p>	<p><b>Population:</b> TBI; <i>Streamed Group (n=65)</i>: Mean Age=44.02yr; Gender: Male=49, Female=16. <i>Control (n=65)</i>: Mean Age=44.46yr; Gender: Male=40, Female=25. <b>Intervention:</b> Retrospective review of patients who received treatment according to either a traditional neurorehabilitation model or a functionally based streamed model. In the streamed model patients were divided into a neurocognitive (NC) stream for predominantly cognitive/behavioural deficits, or a neurophysical (NP) stream for predominantly physical deficits. <b>Outcome Measure:</b> Length of stay, Functional Independence Measure (FIM), and Disability Rating Scale.</p>	<ol style="list-style-type: none"> <li>1. The NP group had higher FIM motor subscale efficiency than controls (0.41 versus 0.29; p=0.01).</li> <li>2. The NC group had less disability (Disability Rating Scale) compared to controls (3.63 versus 5.05, p=0.01).</li> <li>3. Groups did not differ significantly on length of stay, FIM discharge scores, total FIM gain, or total FIM efficiency.</li> </ol>
<p><a href="#">Semlyen et al.</a> (1998) UK PCT N=51</p>	<p><b>Population:</b> TBI; <i>Group 1 (n=33)</i>: Mean Age=36yr; Gender: Male=28, Female=5; Mean Time Post Injury=49.37 days; Mean GCS=5.03. <i>Group 2 (n=18)</i>: Mean Age=30yr; Gender: Male=15, Female=3; Mean Time Post Injury=17.94 days; Mean GCS=5.39. <b>Intervention:</b> Two groups were observed. Group 1 received a coordinated multidisciplinary rehabilitation program in a regional rehabilitation center and Group 2 received single-discipline rehabilitation provided in local hospitals. Assessments were completed at 4wk, 8wk, 12wk, 6mo, 12mo and 24mo post-treatment. Caregivers were assessed at 12wk, 6mo and 12mo after patient’s admission. <b>Outcome Measure:</b> Functional Independence Measure (FIM), Barthel Index (BI), Newcastle Independence Assessment Form (NIAF), and General Health Questionnaire.</p>	<ol style="list-style-type: none"> <li>1. Within the first 6mo of treatment, Group 1 made rapid gains on the BI (p&lt;0.001), NIAF (p&lt;0.0001), FIM Motor (p&lt;0.01) and FIM Cognitive (p&lt;0.05). Group 2 made improvements only up to 12wk post injury on the NIAF (p&lt;0.05) and FIM Cognitive (p&lt;0.05), but not on BI or FIM motor.</li> <li>2. Group 1 demonstrated functional gains over the 12 to 24mo period, making significant gains on the BI and the NIAF (both p&lt;0.05).</li> <li>3. Caregivers of those in the Group 2 reported significantly higher levels of distress for somatic symptoms (p=0.001). Social dysfunction, whilst non-significant, was also much higher for the Group 2 caregivers (p=0.057).</li> </ol>

## Discussion

In a case control study, Cullen et al. (2013) assessed the efficacy of a functionally based neurorehabilitation program that places patients into treatment groups based on their predominant functional deficits. Patients with predominantly physical or cognitive deficits were streamed into a neurophysical or neurocognitive rehabilitation group. When compared with historical controls who received traditional ABI inpatient care, the neurophysical stream had significantly higher functional independence measure efficiency, and the neurocognitive stream had significantly lower disability rating scale scores. It may be more beneficial to rehabilitate patients based on their specific functional or cognitive deficits rather than focusing on general all-inclusive ABI care strategies, or “rehabilitation by diagnosis”.

Another issue is that brain injury often leads to multiple deficits that span across many disciplines. Thus, patients would likely benefit from receiving comprehensive care across multiple areas of expertise.

Semlyen et al. (1998) compared two distinct TBI inpatient rehabilitation strategies. One group of patients received a coordinated, multidisciplinary regional rehabilitation service, and another received a local, single discipline approach. The multidisciplinary rehabilitation group made significant gains up to 6 months according to the functional independence measure, and up to 24 months according to the Barthel Index and Newcastle Independence Assessment Form, whereas the single discipline approach group made significant gains only up to 12 weeks post injury. Notably, the site where the multidisciplinary approach group was treated likely had greater access to resources than other sites, which may have confounded results.

## Conclusions

*There is level 3 evidence that functionally-based streamed models of inpatient ABI care, particularly neurophysical and neurocognitive streams, may improve functional independence measure efficiency and reduce disability rating scale scores, respectively, compared to traditional inpatient rehabilitation.*

*There is level 2 evidence that a coordinated, multidisciplinary inpatient rehabilitation approach may increase the extent of improvement seen in TBI motor and cognitive outcomes as well as lead to longer-term maintenance (up to 24 months) of the treatment effect, compared to single discipline treatment.*



### KEY POINTS

- Functionally-based streamed models of inpatient rehabilitation may improve targeted deficits more than all-encompassing traditional inpatient rehabilitation methods in patients with TBI.
- Compared to a single discipline approach, a coordinated, multidisciplinary approach to inpatient rehabilitation may result in greater functional improvements that are sustained for longer in patients with TBI.

## Intensity of Inpatient Rehabilitation

While patients are undergoing rehabilitation the intensity of therapy provided to them is potentially an important factor in promoting neurological and functional recovery. We review the evidence for increased intensity in this section.

**TABLE 7 | Intensity of Inpatient Rehabilitation Post ABI**

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Hart et al.</a> (2016) Denmark PCT N=274</p>	<p><b>Population:</b> TBI; <i>US Group (n=145):</i> Mean Age=37.5yr; Gender: Male=101, Female=44. <i>Denmark Group (n=129):</i> Mean Age=39.6yr; Gender: Male=103, Female=26. <b>Intervention:</b> Patient outcomes from 2 TBI treatment centers were compared; a US center and a Denmark center where patients received significantly greater intensity and duration of rehabilitation. <b>Outcome Measure:</b> Functional Independence Measure (FIM), Glasgow Outcome Scale (GOS), Disability Rating Scale, Participation Assessment with Recombined Tools, Perceived Quality of Life, Medical Outcome Study 12-Item Short-Form Health Survey, Brief Symptom inventory.</p>	<ol style="list-style-type: none"> <li>1. Injury severity on admission was greater at the Denmark site compared to the US site. The DK center also provided significantly more rehabilitation services for both functional and emotional components (<math>p&lt;0.001</math>).</li> <li>2. After adjusting for injury severity upon admission, there were no significant differences in functional or emotional outcomes between the Denmark and US site at 12 months post-TBI.</li> </ol>
<p><a href="#">Zhu et al.</a> (2001) Hong Kong RCT PEDro=8 N=36</p>	<p><b>Population:</b> TBI; <i>Conventional (n=21):</i> Mean Age=33yr; Gender: Male=17, Female=4; Severity; Severe=13, Moderate=8. <i>Intensive (n=15):</i> Mean Age=30yr; Gender: Male=11, Female=4; Severity: Severe=10, Moderate=5. <b>Intervention:</b> Patients were randomized into two groups: the intensive group received 4hr/day of therapy 5 days/wk while the conventional treatment group received 2 hr/day. Assessments were completed monthly for the first 6mo, then bi-monthly up to 1 yr. <b>Outcome Measure:</b> Glasgow Outcome Scale (GOS) and Functional Independence Measure (FIM).</p>	<ol style="list-style-type: none"> <li>1. A greater proportion of patients from the intensive group achieved “good” GOS outcome 2mo into treatment, compared with the conventional group (40% versus 10%; <math>p=0.046</math>). This difference, however, diminished from 3mo onwards.</li> <li>2. While there were no significant differences in mean FIM motor, cognitive and total scores, there appeared to be a trend in favour of the Intensive group.</li> </ol>
<p><a href="#">Shiel et al.</a> (2001) UK RCT PEDro=7 N=56</p>	<p><b>Population:</b> ABI; <i>Group 1 (n=12):</i> Mean Age=34.2yr. <i>Group 2 (n=13):</i> Mean Age=36.2yr. <i>Group 3 (n=12):</i> Mean Age=37yr. <i>Group 4 (n=14):</i> Mean Age=39yr. <b>Intervention:</b> Patients were randomly assigned to an intervention group with increased therapy intensity or a control group at two separate hospitals. Groups 1 and 2 were recruited from Southampton General Hospital and Groups 3 and 4 were recruited from Poole Hospital. Groups 1 and 3 received intensive therapy and Groups 2 and 4 received routine therapy. <b>Outcome Measure:</b> Functional Independence Measure and Functional Assessment Measure, Length of Stay</p>	<ol style="list-style-type: none"> <li>1. Patients in Groups 3 and 4 were discharged significantly earlier (<math>p=0.004</math>) and received more routine therapy per week (<math>p=0.0099</math>) than patients in Groups 1 and 2.</li> <li>2. Length of stay was not significantly different between the overall intervention group (groups 1 and 3) and the control (groups 2 and 4).</li> <li>3. Patients in Groups 1 and 3 made significantly faster gains in self-care, psychosocial function (both <math>p&lt;0.001</math>), continence (<math>p=0.001</math>), transfers (<math>p=0.002</math>), locomotion, cognition (both <math>p=0.008</math>) and communication (<math>p=0.01</math>) compared to patients in groups 2 and 4.</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Webb &amp; Glueckauf</a> , (1994) USA RCT PEDro=5 N=16	<p><b>Population:</b> TBI; Mean Age=27.4yr; Gender: Male=14, Female=2.</p> <p><b>Intervention:</b> Patients were assigned to either a high (n=8) or low (n=8) involvement neurorehabilitation goal-setting group. Assessments were completed 1wk pre-intervention, 1wk post-intervention and at 2mo follow-up.</p> <p><b>Outcome Measure:</b> Goal Attainment Scaling, Galveston Orientation and Amnesia Test.</p>	<ol style="list-style-type: none"> <li>Both groups made significant improvements in obtaining their goals from pre- to post-intervention (<math>p&lt;0.01</math>) but there were no significant differences between groups.</li> <li>Patients who had high involvement in their neurorehabilitation goal setting maintained their improvements at 2mo follow-up (<math>p&lt;0.001</math>) whereas patients who received low involvement demonstrated a decline in the number of goals attained.</li> </ol>
<a href="#">Cicerone et al.</a> (2004) USA PCT N=56	<p><b>Population:</b> TBI; <i>Group 1 (n=27):</i> Mean Age=37.8yr; Gender: Male=17, Female=10; Mean Time Post Injury=33.9 mo.</p> <p><i>Group 2 (n=29):</i> Mean Age=37.1yr; Gender: Male=23, Female=6; Mean Time Post Injury=4.8 mo.</p> <p><b>Intervention:</b> Patients participated in one of two groups: Group 1 took part in an intensive cognitive rehabilitation program and group 2 was given a standard neurorehabilitation program for 4 mo.</p> <p><b>Outcome Measure:</b> Community Integration Questionnaire (CIQ), Trail-Making Test Parts A and B, California Verbal Learning Test, Controlled Oral Word Association Test, Rey Complex Figure, and Category Test.</p>	<ol style="list-style-type: none"> <li>Both groups showed significant improvements on total CIQ scores following treatment (<math>p&lt;0.001</math>). There was also a between-group difference, with participants in Group 1 showing greater improvement than those in Group 2 (<math>p=0.021</math>).</li> <li>Patients in Group 1 demonstrated significant improvements in neuropsychological function from pre- to post-treatment (<math>p&lt;0.001</math>). Neuropsychological function was not evaluated for Group 2.</li> <li>There was a significant difference in patients' satisfaction with community functioning; patients in Group 2 indicated greater levels of satisfaction than patients in Group 1 (<math>p=0.03</math>).</li> </ol>
<a href="#">Cifu et al.</a> (2003) USA PCT N=491	<p><b>Population:</b> TBI; Mean Age=34.3yr; Gender: Male=354, Female=137; Mean GCS=7.98.</p> <p><b>Intervention:</b> Inpatient rehabilitation data from three medical centers was collected from the Traumatic Brain Injury Model Systems database. Patients received a combination of rehabilitation therapies, including speech, psychological, occupational, and physical therapy.</p> <p><b>Outcome Measure:</b> Functional Independence Measure, Length of Stay (LOS), and hours of therapy.</p>	<ol style="list-style-type: none"> <li>Rehabilitation intensity predicted motor functioning at discharge (<math>p&lt;0.001</math>) but did not predict cognitive gain.</li> <li>Cognition and motor abilities at admission significantly predicted LOS (<math>p&lt;0.01</math>).</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Spivack et al.</a> (1992) USA Case Series N=95	<p><b>Population:</b> TBI; Mean Age=38.6yr; Gender: Male=61, Female=34; Mean Time Post Injury=62.4 days; Mean GCS=8.8.</p> <p><b>Intervention:</b> Data was obtained from monthly medical records at a comprehensive inpatient rehabilitation program. Time spent and intensity of rehabilitation therapies were also observed including physical therapy, cognitive remediation, neuropsychology, and therapeutic recreation.</p> <p><b>Outcome Measure:</b> Physical Performance*, Higher-level Cognitive Skills*, Cognitively Mediated Physical Skills*, Rancho Los Amigos Levels of Cognitive Functioning Scale (RLAS); *Denotes seven-point functional status scales developed by the clinicians within each rehabilitation discipline.</p>	<ol style="list-style-type: none"> <li>1. For physical performance, higher-level cognitive skills and cognitively mediated physical skills, all participants showed improvements from admission to discharge. The short LOS group was significantly better than the long LOS group at admission (<math>p&lt;0.05</math>); however, at discharge all outcome measures were comparable between the 2 groups.</li> <li>2. The effect of training intensity in the first month on physical performance and cognitively mediated physical skills was not significant and approached significance for higher level cognitive skills (<math>p=0.06</math>).</li> <li>3. A significant interaction between training intensity for the first month and RLAS was obtained, with those in the high intensity group showing greater improvements (<math>p&lt;0.05</math>).</li> </ol>
<a href="#">Blackerby,</a> (1990) USA Case Control N=86	<p><b>Population:</b> TBI=83, Other=3.</p> <p><b>Intervention:</b> Retrospective analysis of 2 hospitals that increased their rehabilitation intensity from 5hr/day to 8hr/day, 7 days/wk. Patients who underwent rehabilitation before the intensity increase were compared to those who underwent rehabilitation after.</p> <p><b>Outcome Measure:</b> Length of Stay (LOS).</p>	<ol style="list-style-type: none"> <li>1. Increased rehabilitation therapy resulted in a 31% decrease in LOS for both coma and acute groups at both hospitals in the study (<math>p&lt;0.05</math>).</li> <li>2. Patients in the coma group experienced an average LOS reduction of 48.43 days and the acute group averaged 52.87 days in reduced LOS.</li> </ol>

## Discussion

A number of studies evaluated the efficacy of increased intensity of inpatient rehabilitation on patient outcomes and length of stay following brain injury. In a multicenter PCT, Cifu et al. (2003) examined the efficacy of rehabilitation intensity on functional outcomes at discharge. Rehabilitation intensity predicted motor functioning at discharge but not cognitive gain. Cicerone et al. (2004) found that intensive and structured cognitive rehabilitation therapy (group and individual) and standard neurorehabilitation therapy both resulted in improvements on the community integration questionnaire, however, in the more intensive program, participants made greater gains in cognitive functioning. In another study, Spivack et al. (1992) found that intensity of treatment in the first month of inpatient rehabilitation did not have a significant effect on any of the outcomes measured. However, the authors used functional status scales that were developed by the clinicians within each rehabilitation discipline, and the validity of these scales remains unknown. In a RCT by Zhu et al. (2001), patients were randomized into two groups based on rehabilitation intensity. One group received 4 hours per day of therapy, and another received 2 hours per day, with both groups receiving therapy 5 days per week. The authors reported that significantly more patients from the intensive group achieved good outcomes at 2 months as defined by the Glasgow Outcome Scale, however, this effect was not sustained at 3 months as the conventional therapy group caught up. Moreover, there were no differences between groups in the Functional Independence Measure scores. This study suggests that more intensive rehabilitation may

provide added benefits in the first two to three months post injury, although as time progresses, those who receive less intensive therapy eventually catch up.

Length of stay is another important factor to consider that is closely related to rehabilitation intensity. Blackerby et al. (1990) assessed the effect of different levels of rehabilitation intensity on length of stay. In two separate hospitals, rehabilitation intensity was increased from 5 hours per day to 8 hours per day, 7 days per week. When comparing those who received less intensive therapy with those who received the more intensive regimen, the latter group was discharged an average of 1.5 months earlier. Moreover, the variability in length of stays was decreased in the intensive rehabilitation group, making length of stay easier to predict.

Another factor to consider is the type of rehabilitation that patients are undergoing. Some rehabilitation efforts will benefit more from increased intensity than others. For example, Webb and Glueckauf (1994) found that patients who had greater involvement in goal-setting maintained their improvements at study follow-up. In contrast, those with low involvement in their goal setting showed a decline in the number of goals attained. Not surprisingly, it is beneficial to have a more intense patient involvement in this case, as it ensures the goals are meaningful, and thus the motivation of the patient is increased.

It could also be true that the effectiveness of increased rehabilitation intensity likely depends on the site where the patient is receiving care. If the rehabilitation institution has limited access to resources, this may reduce the efficacy of higher intensity rehabilitation. In a 2-center RCT by Shiel et al. (2001), patients in the intervention group received more intense therapy through the addition of a health care professional (a rehabilitation nurse at one center and an occupational therapist at the other) who provided these extra services as requested by the team. The authors found that, in all domains of the FIM and FAM, the intervention group showed accelerated functional progress when compared to the control group. Interestingly, this did not translate into a shorter length of stay for the intervention group. This may have been related to the long length of stay that was observed in the Southampton intervention group. Length of stay was significantly shorter for both the routine and intervention groups treated at Poole. Importantly, as a result of staffing levels, the mean amount of therapy provided to both subgroups at Poole was significantly greater than that at Southampton. In another study, two rehabilitation sites, one in Denmark and the other in the USA, were compared (Hart et al., 2016). While the Denmark site provided significantly more rehabilitation services for functional and emotional components, no significant differences in functional or emotional outcomes were found at 12 months post TBI. However, this study tracked both inpatient and outpatient rehabilitation services for 1 year post TBI, thus the effect of inpatient rehabilitation intensity alone could not be determined. It is also worth noting that although controlled for in the analysis, the Denmark group had significantly higher injury severity upon admission.

Relatively few studies have evaluated the impact of intensity on rehabilitation, and of the studies that have, results are varied and unclear. Intuitively, it seems reasonable to assume that more intensive therapy will result in more rapid and ultimately greater improvement in recovery from brain injury. Based on the available literature, greater intensity appears to result in a faster recovery in the short term



and therefore shorter lengths of stay, but not necessarily better long-term outcomes, as those with a lower intensity rehabilitation regimen tend to catch up.

## Conclusions

*There is level 1b evidence that more intensive inpatient rehabilitation may improve Glasgow Outcome Scale scores at 2 months, but not necessarily at 3 months and beyond, compared to conventional treatment in patients with TBI.*

*There is level 3 evidence that increasing inpatient rehabilitation intensity may reduce hospital length of stay compared to conventional therapy post ABI.*

*There is level 2 evidence that inpatient therapy intensity post ABI predicts motor functioning at discharge.*

*There is level 2 evidence that a high-level of involvement in neurorehabilitation goal setting may result in a greater number of attained goals being maintained at follow-up (two months) compared to neurobehavioral therapy in individuals with an ABI.*

### KEY POINTS

- Increasing inpatient rehabilitation intensity can reduce hospital length of stay post ABI.
- Increasing inpatient rehabilitation intensity compared to standard therapy, can improve Glasgow Outcome Scale scores and functional outcomes post ABI in the short term.
- The efficacy of increased inpatient rehabilitation intensity post ABI can change based on the rehabilitation institution and available resources.
- Inpatient therapy intensity predicts motor functioning post ABI at discharge.
- High-level involvement in neurorehabilitation goal setting may result in a greater number of attained goals being maintained at follow-up (two months) in individuals with an ABI.

## Timing of Inpatient Rehabilitation

It has long been identified that early onset of therapeutic interventions for those who have sustained a TBI is beneficial. Understanding the ideal time to initiate rehabilitation will help maximize the appropriate use of resources that are available to patients for a limited amount of time. At one end of the spectrum, a comatose patient may be unable to engage in therapy, while at the other end of the spectrum, someone who has made a good recovery has no need for intervention. Several studies have shown that introducing a rehabilitation program during the acute phase assists in the overall recovery of individuals with a TBI (Heinemann AW, 1990). A review by Cope (1995) concluded that those who receive early intervention do in fact have better outcomes than those who do not. Further, León-Carrión et al. (2013) reported that patients who received neurorehabilitation earlier demonstrated better global functioning at discharge than patients who began treatment at a later point. We attempt to address the question of the ideal time to start the rigors of therapy, as well as review the available evidence regarding the effectiveness of delayed ABI rehabilitation in order to maximize patient function and quality of life.



**TABLE 8 |** Timing of Inpatient Rehabilitation Post ABI

Author Year Country Research Design PEDro Sample Size	Methods	Outcomes
<p><a href="#">Formisano et al.</a> (2017) Italy Case Control N=651</p>	<p><b>Population:</b> TBI; Mean Age=43.67yr; Gender: Male=516, Female=135; Severity: Severe. <b>Treatment:</b> Participants were recruited from an inpatient rehabilitation centre and categorized by time from injury to rehabilitation (latency). <b>Outcome Measures:</b> Length of stay (LOS), Disability Rating Scale (DRS).</p>	<ol style="list-style-type: none"> <li>1. There was a significant positive correlation between latency and LOS (p&lt;0.01).</li> <li>2. There was a significant positive correlation between latency and mean admission DRS (p&lt;0.01).</li> <li>3. There was a significant positive correlation between latency and mean discharge DRS (p&lt;0.01).</li> <li>4. There was a positive correlation between latency and the number of participants retransferred to acute care (p&gt;0.05).</li> </ol>
<p><a href="#">Bender et al.</a> (2014) Germany Case Series N=125</p>	<p><b>Population:</b> TBI=38, Intracerebral Hemorrhage=23, Stroke=23, Anoxic Encephalopathy=20, Unknown=1; Mean Age=50.4yr; Gender: Male=73 Female=53. <b>Intervention:</b> Retrospective analysis of a group of patients with severe ABI who participated in an early rehabilitation program (ERP), followed by an inpatient interval rehabilitation program (IRP) a mean of 1.5 years later. <b>Outcome Measure:</b> Goal Attainment Scale, Barthel Index (BI), Functional Independence Measure (FIM), and Coma Remission Scale.</p>	<ol style="list-style-type: none"> <li>1. Thirty-seven percent of IRP inpatients were successful overall in achieving their goals; success rates varied based on primary goals: 86.7% for decannulation, 34.6% for improvements in Activities of Daily Living (ADL), 30% for improvement in dysphagia, 17% for other individual goals (p&lt;0.001).</li> <li>2. Improvement in FIM scores was found during ERP, community care and also IRP (p&lt;0.001).</li> <li>3. BI scores improved significantly during ERP (p&lt;0.001) and continued to improve during IRP (p&lt;0.001).</li> </ol>
<p><a href="#">High et al.</a> (2006) USA PCT N<sub>Initial</sub>=167, N<sub>Final</sub>=141</p>	<p><b>Population:</b> TBI; <i>Group 1 (n=115):</i> Mean Age=31.5yr; Gender: Male=86, Female=29. <i>Group 2 (n=23):</i> Mean Age=32.8yr; Gender: Male=14, Female=9. <i>Group 3 (n=29):</i> Mean Age=27.2yr; Gender: Male=18, Female=11. <b>Treatment:</b> Patients were enrolled in a comprehensive, integrated post-acute brain injury rehabilitation program. Patients were grouped depending on length of time between injury and admission: &lt;6mo (Group 1), 6-12mo (Group 2), and &gt;12mo (Group 3). Patients participated in an interview at admission, discharge and at approximately 1.5yr follow-up. <b>Outcome Measure:</b> Disability Rating Scale (DRS), Supervision Rating Scale (SRS), and Community Integration Questionnaire (CIQ).</p>	<ol style="list-style-type: none"> <li>1. For those in Group 1, DRS scores from admission to discharge improved significantly (p&lt;0.001). Such improvements were not seen in Groups 2 or 3.</li> <li>2. SRS scores decreased from admission to follow-up for Group 1 at all time-points (all p&lt;0.001).</li> <li>3. Groups 2 and 3 reported significant decrease in supervision between admission and discharge (p=0.001 and p=0.002 respectively) but no significant change was observed between discharge and follow-up.</li> <li>4. All groups demonstrated improvements in CIQ between admission and discharge (p&lt;0.001) and between discharge and follow-up (p=0.003).</li> <li>5. Social integration improved significantly between admission and follow-up (p=0.014) for all groups.</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcomes
<p><a href="#">Wagner et al.</a> (2003) USA Case Control N=1,866</p>	<p><b>Population:</b> TBI; <i>Group 1 (n=520):</i> Mean Age=48.5yr; Gender: Male=339, Female=181; Severity: Severe=178, Mild/Moderate=326. <i>Group 2 (n=1,346):</i> Mean Age=39.4yr; Gender: Male=939, Female=407; Severity: Severe=90, Mild/Moderate=1163. <b>Treatment:</b> A comparison was conducted between Group 1, patients who received physical medicine and rehabilitation consultation and those who did not (Group 2). Data was extracted from hospital records. <b>Outcome Measure:</b> Modified Functional Independence Measure (mFIM).</p>	<ol style="list-style-type: none"> <li>1. Patients in Group 1, compared to Group 2, were more likely to have worse mFIM scores at acute discharge (p=0.05), have at least one premorbid condition (p=0.002) and have significantly longer length of stay (11.85 days versus 2.47 days, p&lt;0.001).</li> <li>2. For group 1, when the consultations occurred earlier (&lt;48hr after hospital admission) patients experienced significantly better mFIM scores for transfers and locomotion (both p=0.05) and had significantly shorter acute length of stay (p=0.001).</li> </ol>
<p><a href="#">Edwards et al.</a> (2003) USA Cohort N=290</p>	<p><b>Population:</b> ABI: TBI=110, Intracerebral Hemorrhage/Subarachnoid Hemorrhage/Cerebral Infarction=122, Other=58; Mean Age=38yr; Gender: Male=193, Female=97. <b>Treatment:</b> Data was extracted from a hospital database. Patient assessments were conducted within 4wk of admission, every 6-8wk and at discharge. Patients were retrospectively split into two group, those admitted &lt;200 days post-injury (n=264) and those admitted &gt;200 days post-injury (n=26). <b>Outcome Measure:</b> Length of Stay, Barthel Index (BI), and Functional Independence Measure (FIM).</p>	<ol style="list-style-type: none"> <li>1. Rehabilitation length of stay was similar for the two groups.</li> <li>2. Lower BI and FIM scores at admission were significant predictors for increased length of stay for all patients (both p&lt;0.001).</li> <li>3. Discharge BI and FIM scores were lower in the admitted &gt;200 days post-injury group than the &lt;200 days post-injury group (BI, 11 versus 14; FIM, 77 versus 92 respectively), but the differences were not significant.</li> </ol>
<p><a href="#">Tuel et al.</a> (1992) USA Case Series N=49</p>	<p><b>Population:</b> ABI; Mean Age=23.6yr; Gender: Male=38, Female=11; Mean Time Post Injury=2.9yr. <b>Intervention:</b> Data was obtained from records of patients readmitted to inpatient rehabilitation more than 12mo after injury. <b>Outcome Measure:</b> Barthel Index (BI).</p>	<ol style="list-style-type: none"> <li>1. Fifty-three percent (n=26) showed improvement (mean BI gain of 11.2 points).</li> <li>2. Statistically significant improvements of BI scores were shown from re-admission to discharge (p=0.0001).</li> <li>3. Length of readmission was significantly correlated with improvements in BI (p=0.0016).</li> </ol>
<p><a href="#">Cope and Hall,</a> (1982) USA Case Control N=36</p>	<p><b>Population:</b> ABI; <i>Early Group (n=16):</i> Mean Age=29yr; Gender: Male=9, Female=7; Mean Time Post Injury=20.88 days; Mean GCS=5.54. <i>Late Group (n=20):</i> Mean Age=29.15yr; Gender: Male=15, Female=5; Mean Time Post Injury=61.35 days; Mean GCS=5.11. <b>Treatment:</b> Patients were retrospectively assigned to one of two groups: an Early Rehabilitation Group which consisted of patients admitted to a rehabilitation facility at ≤35d post-injury or a Late Rehabilitation Group with patients admitted to a rehabilitation facility at &gt;35d post-injury. <b>Outcome Measure:</b> Disability Rating Scale, Glasgow Outcome Scale, Social Status Outcome (SSO).</p>	<ol style="list-style-type: none"> <li>1. Both groups reached equivalent levels of functional recovery at discharge and SSO ratings at 2yr post-injury.</li> <li>2. Those in the Late Group spent significantly more time in acute care (p=0.001) and inpatient rehabilitation (p=0.01) than the Early Group.</li> <li>3. At 2mo post-injury, patients in the Early Group experienced significantly less psychological impairment (p=0.02), and fewer problems with bowel and bladder function (p=0.05) than the Late Group.</li> </ol>

## Discussion

In this section, our analysis is two-fold. First, we review studies that evaluate the effect of earlier acute intervention on functional outcomes and length of stay, then we assess the relative efficacy of rehabilitation in the chronic phase post ABI.

In a case control study by Formisano et al. (2017), patients from an inpatient rehabilitation center were categorized by time from injury to initiation of rehabilitation. Patients who began rehabilitation sooner after injury had lower length of stays, lower initial disability rating scale scores, and higher mean discharge disability rating scale scores. Wagner et al. (2003) examined the proper timing for physical medicine and rehabilitation consultation. Using multivariate analysis, the authors found that when physical medicine and rehabilitation consultations occurred earlier (<48 hours after hospital admission) patients experienced significantly better Functional Independence Measure scores for transfers and locomotion and had significantly shorter lengths of stay. Cope and Hall (1982) reported that those in a late intervention (>35 days) group spent significantly more time in acute care and inpatient rehabilitation. Edwards et al. (2003) compared 26 patients admitted to inpatient rehabilitation more than 200 days after injury to 264 patients admitted to inpatient rehabilitation less than 200 days after injury. Although it was not significant, the discharge scores on the Barthel Index and Functional Independence Measure were lower in the former group than in the latter. Rehabilitation LOS was also similar for the two groups. Generally, it seems that patients who sustain an ABI benefit from earlier initiation of rehabilitation.

However, a number of studies have also shown that although it is beneficial to begin rehabilitation soon after sustaining an ABI, rehabilitation efforts in the chronic phase can still result in significant improvements. Bender et al. (2014) reported an improvement in Functional Independence Measure scores during early rehabilitation, community care, and inpatient interval rehabilitation. Although patients entered the inpatient interval rehabilitation program an average of 1.5 years after discharge from the early rehabilitation program, they still demonstrated improvement-rate increases comparable to initial rehabilitation levels, where the greatest gains are said to be made, highlighting the benefit of additional rehabilitation at later stages of recovery. This point has been made by earlier studies as well. A study noted that 53% of patients readmitted to inpatient rehabilitation at more than 12 months post injury showed statistically significant improvement on Barthel Index scores from readmission to discharge (Tuel et al., 1992). In a PCT, modest findings were reported from High et al. (2006). All three-time groups (time since injury of less than 6 months, 6 to 12 months, greater than 12 months) demonstrated a significant decrease in required supervision from admission to discharge; however, the less than six-month group continued to improve through to follow-up. Similarly, Rosenbaum et al. (2018) observed that individuals with moderate-severe TBI continued to make significant functional gains when provided with long-term inpatient interdisciplinary rehabilitation (2.6-3.15 hours per day 5 days per week), regardless of age, injury severity, or time post injury.

### Conclusions

*There is level 2 evidence that inpatient rehabilitation within 35 days post ABI is associated with better outcomes such as shorter comas and hospital length of stay, higher cognitive levels at discharge, better Functional Independence Measure scores, and a greater likelihood of discharge to home, compared to rehabilitation initiated after 35 days post-ABI.*


*There is level 2 evidence that individuals with an ABI can still benefit from inpatient rehabilitation efforts initiated more than 12 months after sustaining an ABI.*



#### KEY POINTS

- Early inpatient rehabilitation is associated with better outcomes in individuals post ABI.
- Inpatient rehabilitation in the chronic phase of ABI can still yield meaningful results.

## Outpatient Rehabilitation and Community Reintegration



Outpatient rehabilitation is often the least organized branch of ABI care. Patients discharged home often receive no therapy or minimal support depending on their level of need and payment status. In a well-structured outpatient facility in Canada, patients typically attend therapy two to three times per week and have access to an occupational therapist, physiotherapist, speech language pathologist, social worker, physiatrist, neuropsychologist, and neuropsychiatrist (Cullen, 2007). At some facilities, for example in Hamilton, Ontario, patients also receive the services of a rehabilitation counselor. Access to programs like these often depends on funding. Patients with private insurance from motor vehicle accidents are 1.6 times more likely to be discharged home with supportive services than those without (Kim et al., 2006). Drag et al. (2013) reported that US TBI-veterans were significantly more likely to utilize outpatient services and were almost nine times more likely to be hospitalized than non-TBI veterans. The authors argue that earlier intervention and increased monitoring may be needed to reduce the burden on outpatient healthcare (Drag et al., 2013). In a study by Leith et al. (2004), focus groups of patients and families were questioned regarding their perceived post-discharge needs. The areas of need included: early, continuous, comprehensive service delivery, information and education, formal and

informal advocacy, empowerment of persons with TBI and their families, and human connectedness and social belonging (Leith et al., 2004). Subsequently, a survey was conducted in the US to identify the availability of community information resources post-ABI (Sample & Langlois, 2005). Three recommendations for improvement were made: expand the population targeted for linkage to services, improve access to information about available services, and increase the availability of services (Sample & Langlois, 2005).

Residential care facilities are generally not-for-profit, government sponsored agencies that offer access to support in a secure environment with staff specifically trained in ABI care. Resources often include rehabilitation therapists, behaviour therapists, social workers, and case managers, with supervision by certified psychologists (Powell et al., 2002). These facilities aim to allow patients with ABI an extended system of support, with opportunities for long-term rehabilitation. However, they are generally expensive, and access is often limited by the patient's ability to pay for care. Alternatives include hospital based outpatient facilities where patients drop in several times a week for care (Cullen, 2007) or mobile rehabilitation teams which visit the patient in their home (Ponsford et al., 2006). Programs targeting specific goals including social interaction (Cope et al., 2005), driving (Rapport et al., 2008) and competitive employment (Willer et al., 1999) also exist. They generally take place on a one-to-one basis in home or in the community and patients often rate these final steps as the most important in returning to normalcy (Evans, 1997).

While most patients move back to their former living environment with therapy intervention provided for them in the home or community, some go on to other facilities that may provide longer duration treatment for those that are slow-to-recover. There are numerous models of care within community-based rehabilitation. Regarding social re-entry, Martelli (2012) argues that peer support interventions are effective in reintegrating individuals and should be incorporated into community rehabilitation models. Avocational and vocational reintegration are also important factors as the former can improve functionality, strength, and endurance, while the latter improves productivity, self-worth, and life satisfaction. Vocational reintegration can be facilitated through the case coordination model whereby the patient collaborates with a case coordinator who assesses the services needed and makes appropriate referrals on the patient's behalf (Martelli et al., 2012). Studies discussing a wider range of vocational rehabilitation interventions can be found in Module 13 (Community Reintegration). Lack of services could be an issue, thus a supported employment program or a program-based model could also be utilized.

Ashley et al. (2012) proposed a new community-based interventional model that focuses on cognitive attributes. The proposed model incorporates tasks that assist with attention, perceptual processing, and categorization. The authors point out that previous research has found re-establishing neuronal connections that become damaged after a TBI leads to greater cognitive functioning in patients. For the model to be successful, Ashley et al. (2012) state that the tasks need to be errorless, with high levels of repetition and redundancy in order for the intervention to be successful. As this model of care can be extremely time-consuming, it is suggested that this be utilized in a community-based rehabilitation center.

**TABLE 9 |** Outpatient Rehabilitation and Community Reintegration Post ABI

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Poncet et al.</a> (2018) France Pre-post N= 7</p>	<p><b>Population:</b> ABI Group (N= 7): Mean Age=38yr (10.1); Gender: Male=43%, Female=57%; Mean time post injury= n/a, at least 2 months discharged to home; GCS= n/a. <b>Intervention:</b> Participants underwent a multidisciplinary ABI outpatient rehabilitation program 5 days a week for 7 weeks. Assessed at pre, post, 3 months, and 6 months after program <b>Outcomes:</b> The Cooking Task, Instrumental Activities of Daily Living Profile (IADL Profile), Assessment of Life Habits (LIFE H), Berg Balance Scale (BBS), Six-minute Walk Test (6MWT), 10m Walk Test (10MWT), Box and Blocks Test (BBT), Montgomery Asberg Depression Rating Scale (MADRS), Hamilton Depression Rating Scale (HDRS), The Evaluation Test of Attention (ETA), Auditory Verbal Learning Test.</p>	<ol style="list-style-type: none"> <li>1. Non-overlap of all pairs (NAP) scores and two standard deviation method (SDB) was used to determine significance.</li> <li>2. Strong improvement was seen on the Cooking Task for 6/7 participants from pre to post, with four showing improvements at 3 and 6 months.</li> <li>3. Strong improvement was seen on the IADL Profile for 6/7 participants, with four improving from post to 6 months.</li> <li>4. Strong improvement was seen on the LIFE H for 4/7 participants from pre to post.</li> <li>5. No improvements seen on other measures.</li> </ol>
<p><a href="#">Peirone et al.</a> (2014) Italy RCT PEDro=8 N=16</p>	<p><b>Population:</b> ABI; <i>Intervention Group (n=8):</i> Mean Age=39.62yr; Gender: Male=4, Female=4; Mean Time Post Injury=14 mo. <i>Control Group (n=8):</i> Mean Age=40.5yr; Gender: Male=5, Female=3; Mean Time Post Injury=14.5 mo. <b>Intervention:</b> All patients received 50 min of physiotherapy 3x/wk for 7 wk. The intervention group also received additional dual-task exercises 6x/wk including balance and body stability whilst performing a motor task (throw/catch a ball) or a cognitive test. <b>Outcome Measure:</b> Balance Evaluation System Test (BEST), Activities-specific Balance Confidence Scale (ABC), and Goal Attainment Scaling (GAS).</p>	<ol style="list-style-type: none"> <li>1. Both the Intervention group and the control group improved significantly on the BEST (p=0.014 and p=0.02 respectively) but when comparing the two, the Intervention group displayed significantly greater improvements (p=0.008).</li> <li>2. On the ABC, the intervention group made significant improvements from baseline (p=0.01). There was no significant difference between the groups at the end of the study.</li> <li>3. Both intervention and control groups experienced significant improvements on the GAS (p=0.02 and p=0.01 respectively) but no significant difference was found between the two groups.</li> </ol>



Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Eicher et al.</a> (2012) USA Cohort N=604</p>	<p><b>Population:</b> ABI; <i>Group 1 (n=235):</i> Mean Age=41.65yr; Gender: Male=136, Female=99. <i>Group 2 (n=78):</i> Mean Age=38.95yr; Gender: Male=62, Female=16. <i>Group 3 (n=246):</i> Mean Age=35.42yr; Gender: Male=185, Female=61. <i>Long-term Group 4 (n=45):</i> Mean Age=35.78yr; Gender: Male=33, Female=12.</p> <p><b>Intervention:</b> Four rehabilitation programs were assessed: Group 1 received Intensive Outpatient &amp; Community-based Rehabilitation, Group 2 received Intensive Residential Rehabilitation, Group 3 received Long-term Residential Supported Living, and Group 4 received Long-term Community-based Supported Living. Mean follow-up time was 5-6mo for Groups 1, 2 and 4, and 8mo for Group 3.</p> <p><b>Outcome Measure:</b> Mayo-Portland Adaptability Inventory (MPAI-4).</p>	<ol style="list-style-type: none"> <li>1. Programs with an intense rehabilitation program (Groups 1 and 2) demonstrated significant improvements on the MPAI-4 compared to Groups 3 and 4 (p=0.002).</li> <li>2. At baseline assessment, patients in Groups 1 and 2 scored significantly lower than patients in Groups 3 and 4 in adjustment (p&lt;0.001) and ability (p&lt;0.05).</li> <li>3. At the second assessment, patients in Groups 1 and 2 scored significantly better on ability (p&lt;0.006) and participation (p&lt;0.001) than patients in Groups 3 and 4.</li> <li>4. There was no statistical difference on adjustment at second assessment.</li> </ol>
<p><a href="#">Braunling-McMorrow et al.</a> (2010) USA PCT N=205</p>	<p><b>Population:</b> TBI; <i>Neurorehabilitation Group (NR; n=129):</i> Mean Age=36.6yr; Gender: Male=89, Female=40; Mean Time Post Injury=15 mo. <i>Neurobehavioural Group (NB; n=76):</i> Mean Age=32yr; Gender: Male=63, Female=13; Mean Time Post Injury=56 mo.</p> <p><b>Intervention:</b> Individuals were divided into 2 groups: those receiving NR services and those requiring specialized NB services due to behavioural or psychiatric issues. Both groups participated in behavioural and cognitive therapy programs. Assessments were completed at pre-admission, admission, discharge, and 3mo, 6mo and 12mo post-discharge.</p> <p><b>Outcome Measure:</b> Functional Area Outcome Menu.</p>	<ol style="list-style-type: none"> <li>1. Individuals in both groups showed significant functional gains from admission to discharge (p&lt;0.001).</li> <li>2. Functional gains were also made from admission to 1yr follow up (p&lt;0.001).</li> <li>3. The NR group made greater overall gains than those in the NB group (p&lt;0.001).</li> <li>4. Gains made by both groups were noted at the 1yr follow up; whereas NB group continued to make significant gains from discharge to follow-up (2.93 to 3.23; p&lt;0.05), the NR group maintained their gains (3.68 to 3.60; p&gt;0.05).</li> <li>5. Results also found that NR patients admitted within 6mo of injury made the greatest improvement (p&lt;0.001). There was no significant effect of time post injury for the NB group.</li> </ol>
<p><a href="#">Hassett et al.</a> (2009) Australia RCT PEDro=8 N=62</p>	<p><b>Population:</b> TBI; <i>Fitness Group (n=32):</i> Mean Age=35.4yr; Gender: Male=27, Female=5; Severity: Very Severe=9, Extremely Severe=23. <i>Home Group (n=30):</i> Mean Age=33yr; Gender: Male=26, Female=4; Severity: Very Severe=11, Extremely Severe=19.</p> <p><b>Intervention:</b> Patients were randomly assigned to receive either a supervised fitness center-based therapy program or an unsupervised home-based program. Assessments were completed at baseline, at completion of the intervention and at 3mo follow-up.</p> <p><b>Outcome Measure:</b> Goals, 20 m Shuttle Test, Body Mass Index, Waist-to-Hip-ratio, and Sydney Psychosocial Reintegration Scale (SPRS).</p>	<ol style="list-style-type: none"> <li>1. Patients in both groups improved in cardio-respiratory fitness but there were no significant differences between groups.</li> <li>2. At the end of the intervention, the fitness group reported significantly greater total scores on SPRS (p=0.033), but the difference was not significant at follow-up.</li> <li>3. The fitness group also reported significantly more goals achieved at the end of the intervention (p=0.005) but this also became non-significant at follow-up.</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Ponsford et al.</a> (2006) Australia Cohort N=154</p>	<p><b>Population:</b> TBI; <i>Community Group (n=77)</i>: Mean Age=35.43yr; Gender: Male=56, Female=21; Mean GCS=8.22. <i>Outpatient Group (n=77)</i>: Mean Age=33.78yr; Gender: Male=56, Female=21; Mean GCS=7.76. <b>Intervention:</b> Patients treated in a community-based rehabilitation program were matched with patients who attended the hospital for outpatient rehabilitation. Assessments were completed every 3mo during treatment then at 1 and 2yr follow-ups. <b>Outcome Measure:</b> Craig Handicap Assessment and Reporting Technique, and Structured Outcome Questionnaire.</p>	<ol style="list-style-type: none"> <li>1. Patients treated in the community were significantly more dependent on support from close others (p=0.008), less independent in mobility (p=0.005), had greater difficulty with motor speech (p=0.005) and following conversations (p=0.001), and displayed more inappropriate social behaviours (p=0.009) than the outpatient group.</li> <li>2. Patients treated in the community demonstrated increased physical independence (p=0.004) compared to patients in the outpatient group.</li> </ol>
<p><a href="#">Powell et al.</a> (2002) UK RCT PEDro=8 N=94</p>	<p><b>Population:</b> TBI; <i>Outreach Group (n=48)</i>: Mean Age=34yr; Gender: Male=37, Female=11; Mean Time Post Injury=4yr. <i>Information Group (n=46)</i>: Mean Age=35yr; Gender: Male=34, Female=12; Mean Time Post Injury=2.7 yr. <b>Intervention:</b> Patients were randomly allocated to one of two groups. The Outreach group received multi-disciplinary therapy for 2 hr/wk for a mean of 27.3wk in a community setting and the information (control) group received a specially collated booklet with resources. Patients were assessed at 18mo and 40 mo. <b>Outcome Measure:</b> Barthel Index (BI), Functional Independence Measure + Functional Assessment Measure (FIM+FAM), and Brain Injury Community Rehabilitation Outcome-39 (BICRO-39).</p>	<ol style="list-style-type: none"> <li>1. Of the 92 participants who completed the study, 35% of the Outreach group improved their scores on the BI compared to 20% of the Information group (p&lt;0.05).</li> <li>2. Improvements for FIM+FAM scores approached statistical significance when measuring level of personal care (p&lt;0.06) and cognition (p&lt;0.09) for the Outreach group compared to the information group. All other FIM+FAM subscales were non-significant.</li> <li>3. The Outreach group demonstrated significantly greater improvement on the BICRO-39 than the Information group (p&lt;0.05).</li> </ol>
<p><a href="#">Willer et al.</a> (1999) USA/Canada Case Control N=46</p>	<p><b>Population:</b> TBI; <i>Treatment Group (n=23)</i>: Mean Age=33.42yr; Gender: Male= 20, Female=3; Mean Time Post Injury=3.05yr. <i>Control Group (n=23)</i>: Mean Age=34.76yr; Gender: Male=20, Female=3; Mean Time Post-Injury=4.66yr. <b>Intervention:</b> Patients admitted to a community-based residential rehabilitation treatment center were compared to a matched control group receiving rehabilitation at home or in a nursing facility. Assessments were completed at admission, discharge and at 1yr follow-up by staff or relatives. <b>Outcome Measure:</b> Health Activity Limitation Survey and Community Integration Questionnaire (CIQ).</p>	<ol style="list-style-type: none"> <li>1. Patients in the treatment facility showed significant improvement in motor and cognitive functioning at 1yr follow-up compared to the controls (p&lt;0.05).</li> <li>2. The two groups did not differ significantly on the CIQ at discharge or at follow-up, but the treatment group demonstrated significantly greater improvement from admission to discharge than the control group (p&lt;0.001).</li> </ol>

PEDro = Physiotherapy Evidence Database rating scale score (Moseley et al., 2002)

## Discussion

In terms of outpatient and community care, there are several similarities to inpatient rehabilitation. A multidisciplinary approach is still favourable for outpatient services, and timely rehabilitation is imperative as patients are often sent home too early and referred to outpatient services too late (Jeyaraj



et al., 2013; Poncet et al., 2018). Braunling-McMorrow et al. (2010) looked at the benefits of participation in a weekly program that included both behavioural and cognitive therapies that would teach participants to respond to various life events appropriately and allow for greater independence. Those in the neurobehavioural group admitted within the first six months of injury showed greater improvement than those admitted later. The study authors suggest that injury severity may have been a factor, with more severe cases being admitted sooner. As well, for those admitted later, gains had already been made and this may have made the gains in the program appear less significant (Braunling-McMorrow et al., 2010).

Similar to inpatient rehabilitation, intensity of care is an important factor to consider when patients are discharged back into the community and receive care on an outpatient basis. A longer duration of rehabilitation is generally suggested, but with lower intensity to allow patients more time to integrate back into daily life. However, one study has provided evidence that outpatient rehabilitation programs with greater intensity appear to be more effective. Eicher et al. (2012) compared four different outpatient and community rehabilitation programs: an intensive outpatient and community-based program, an intensive residential rehabilitation program, long-term residential supported living, and long-term community-based supported living. The more intensive programs provided more functional benefits, whereas supported living programs resulted in relatively stable scores on the Mayo-Portland Adaptability Inventory. Jeyaraj et al. (2013) note that there is a need to train clinicians who provide community services about how best to assist individuals with ABI and increase the amount of community resources available to them. More studies are required to determine an optimal balance between providing intense outpatient rehabilitation programs versus allowing patients time to reintegrate into society and tackle the responsibilities of daily life in the community.

A number of studies have assessed the effectiveness of different types of outpatient rehabilitation programs. Ponsford et al. (2006) compared outpatients treated in the community to those who returned to the hospital for outpatient care. The findings indicate that patients who received outpatient care were significantly less dependent on support from close others, more independent in mobility, displayed fewer inappropriate social behaviours, and had less difficulty with motor speech and following conversations than those receiving community-based rehabilitation. No significant differences were shown in terms of employment outcomes. One of the main targets for community-based rehabilitation is to assist the patient in their transition into independent living. Independence is a key component of self-efficacy and allows us to live an autonomous life. Powell et al. (2002) randomly assigned patients with TBI to an outpatient support program where patients received two to six hours of therapy a week at home or in another community setting, or to a control group that received an information session at home. Patients in the intervention group showed improvements in cognitive functioning, mobility, and personal wellbeing. Areas such as socializing and competitive employment rates showed no relative difference between groups. The authors suggest that this reflects external influences beyond the control of the rehabilitation team. The authors recommend that this type of outpatient approach be applied to a broader range of patients with ABI in a larger trial to confirm their results. In terms of where outpatient services are provided, one study found that patients with TBI given rehabilitation in a residential treatment center made significantly greater gains in terms of motor and cognitive functioning than those

receiving rehabilitation in a nursing facility or at home (Willer et al., 1999). The groups, however, did not differ at discharge or at a one year follow-up on a measure of community integration (Willer et al., 1999).

Two RCTs specifically looked at the relative effectiveness of different outpatient exercise/physiotherapy programs. Hassett et al. (2009) randomized patients to a supervised fitness center-based exercise program or an unsupervised home-based program and found that both groups at follow-up made comparable gains in terms of psychosocial and physical functioning. Although the fitness center-based group achieved more goals post intervention, the difference was not significant at follow-up. This study highlights that these programs are equally as effective which is positive for individuals who cannot access or prefer not to attend community center fitness programs. In a RCT conducted by Peirone et al. (2014), while all patients received physiotherapy to target balance impairments, those in the intervention group also received a dual-task home-based program. The program was provided six days a week over seven weeks and resulted in this group making significantly greater gains in terms of balance than the control group. While improvements were also shown on the goal attainment scale and a balance confidence scale, the between-group differences were not significant. Unfortunately, the results are hard to interpret due to being underpowered and the inability to distinguish whether the improvements are based on the program itself or simply the increase in rehabilitation intensity as a result of adding an additional therapy.

## Conclusions

*There is level 2 evidence that earlier initiation of outpatient and community-based rehabilitation is associated with better functional outcomes compared to residential rehabilitation in individuals with an ABI.*

*There is level 2 evidence that more intensive outpatient rehabilitation programs are associated with better functional outcomes compared to standard therapy in individuals with an ABI.*

*There is level 2 evidence that compared to individuals with an ABI who are treated in the community, those treated at an outpatient clinic may be less dependent on support from others, more independent in mobility, display fewer inappropriate social behaviours, and have less difficulty with motor speech.*

*There is level 1b evidence that a supervised fitness-center based program may be equally as effective as an unsupervised home-based program for improving cardiorespiratory fitness in individuals with an ABI.*

*There is level 3 evidence that outpatient care provided at a residential treatment center may improve motor and cognitive function to a greater extent than when care is provided at a nursing facility or at home in individuals with a TBI.*



**KEY POINTS**

- Earlier outpatient rehabilitation is associated with better outcomes post ABI.
- More intensive outpatient rehabilitation is associated with better functional outcomes post ABI. However, this may not be the case if intensity is high enough to interfere with a patient’s ability to perform day to day responsibilities.
- Compared to individuals with an ABI who are treated in the community, those treated at an outpatient clinic may be less dependent on support from others, more independent in mobility, display fewer inappropriate social behaviours, and have less difficulty with motor speech.
- Outpatient care provided at a residential treatment center may improve motor and cognitive function to a greater extent than when care is provided at a nursing facility or at home in individuals with a TBI.

## Vocational Rehabilitation

Returning to work following ABI is one of the most challenging tasks that a patient will face in the course of their recovery. The work environment often exposes individuals to physical stressors, cognitive challenges, and emotional strain. However, given the financial burden of not being able to work, for most individuals, it is a very important aspect of full reintegration into society and return to independent living.

**TABLE 10 |** Intensity of Outpatient Rehabilitation for Vocational Rehabilitation Post ABI

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Ownsworth et al.</a> (2008) Australia RCT PEDro=8 N=35	<p><b>Population:</b> TBI=21, Stroke=12, Tumor=2; Mean Age=43.89yr; Gender: Male=19, Female=16; Mean Time Post Injury=5.29 yr.</p> <p><b>Intervention:</b> Patients were randomly allocated to one of six intervention groups: group-based support, individual occupation-based support, or a combination of the two (each of these three interventions had its own waitlist which served as a control group). Relatives and patients completed questionnaires at baseline, post intervention and at 3mo follow-up.</p> <p><b>Outcome Measure:</b> Canadian Occupational Performance Measure (COPM), Brain Injury Community Rehabilitation Outcome-39 (BICRO-39), and Patient Competency Rating Scale (PCRS).</p>	<ol style="list-style-type: none"> <li>1. COPM performance self-ratings improved significantly from pre- to post-assessment following the individual and combined interventions (both <math>p &lt; 0.01</math>), but not the group intervention (<math>p = 0.029</math>). A similar pattern was seen for relatives’ ratings (<math>p &lt; 0.01</math>, <math>p &lt; 0.025</math>, <math>p = 0.028</math>).</li> <li>2. COPM performance self-ratings improved significantly from pre-assessment to 3mo follow-up in the individual (<math>p &lt; 0.025</math>), group and combined (both <math>p &lt; 0.01</math>) intervention groups. A similar pattern was seen for COPM relatives’ ratings (<math>p &lt; 0.01</math>, <math>p &lt; 0.01</math>, <math>p &lt; 0.025</math>).</li> <li>3. Psychological improvements measured by the BICRO-39 were demonstrated from pre- to post-assessment by the patients in the group-based support intervention (<math>p &lt; 0.01</math>) and between pre-assessment and follow-up (<math>p = 0.025</math>). The individual-based support intervention also</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
		<p>demonstrated improvement between pre-assessment and follow-up (<math>p &lt; 0.025</math>).</p> <p>4. Relatives of the patients in the individual-based intervention reported significant improvements on the PCRS from pre- to post-assessment (<math>p &lt; 0.025</math>). Significant improvement was also reported by relatives of group-based patients at 3mo follow-up (<math>p &lt; 0.025</math>).</p>
<p><a href="#">Malec &amp; Degiorgio</a> (2002) USA Secondary Analysis of PCT N=114</p>	<p><b>Population:</b> TBI=73, Other=41; Mean Age=37.4yr; Gender: Male=70, Female=44; Mean Time Post Injury=65.5mo; Severity: Severe=64, Moderate=8, Mild=24, Undetermined=18.</p> <p><b>Intervention:</b> Patients in three rehabilitation pathways were observed: Specialized Vocational Services (SVS) only (n=49), SVS and a 3 hr/wk Community Reintegration Outpatient Group (CROG; n=21), and SVS and a 6 hr/day comprehensive day treatment (CDT; n=44). Assessments were completed by patients and staff at yr follow-up.</p> <p><b>Outcome Measure:</b> Mayo-Portland Adaptability Inventory (MPAI), Vocational Independence Scale, and community-based employment status (CBE).</p>	<ol style="list-style-type: none"> <li>1. No significant differences in CBE rates between all three groups: SVS=77%, CROG=85%, CDT=84% (<math>p &gt; 0.1</math>).</li> <li>2. There were no significant differences between patient MPAI scores across all three groups, except for driving (<math>p &lt; 0.001</math>) with 71% of the SVS group and 60% of the CROG group receiving no driving concerns from others compared to only 26% for the CDT group.</li> <li>3. There were significant differences in staff MPAI scores with the SVS group experiencing fewer memory (<math>p &lt; 0.01</math>) and problem-solving (<math>p &lt; 0.001</math>) impairments, and greater attention and concentration abilities (both <math>p &lt; 0.001</math>) compared to the other groups.</li> <li>4. Patients in the CROG group were more likely to live independently than those in the SVS or CDT groups (<math>p &lt; 0.001</math>).</li> </ol>

### Discussion

Two studies have focused on the ability of outpatient or community rehabilitation to help patients with ABI return to work. Ownsworth et al. (2008) performed a RCT to compare individual, group, and combined intervention formats for goal attainment and psychosocial functioning. Interventions were administered for 3 hours per week for 8 weeks. The individual occupation-based support contributed to gains in performance in goal-specific areas. The combined intervention was associated with maintained gains in satisfaction and performance, while the group and individual interventions were more likely to result in gains in behavioural competency and psychological well-being. These findings provide support for brief intervention formats for individuals in the chronic post-injury phase. In another study, Malec and Degiorgio (2002) reported that patients in three different rehabilitation pathways, who differed in terms of cognitive functioning and disability, were able to succeed in terms of community-based employment. The study highlights the need for an individualized approach to ensure successful integration into the community. The intensity of therapy and the resources and interventions offered must match the individual’s needs, severity of injury, and goals, among other factors (Malec & Degiorgio,

2002).

### Conclusions

There is level 1b evidence that neither individualized nor general vocational rehabilitation programs may improve performance in goal-specific areas compared to waitlist controls in individuals with ABI.

There is level 2 evidence that combining specialized vocational rehabilitation services with a community reintegration outpatient group intervention or comprehensive day treatment may not improve community-based employment compared to specialized vocational rehabilitation alone in individuals with ABI.



### KEY POINTS

- Individualized and group vocational rehabilitation programs can improve goal-specific performance and behavioural competency/psychological well-being in individuals post ABI, respectively.
- Combining specialized vocational rehabilitation services with a community reintegration outpatient group intervention or comprehensive day treatment may not improve community-based employment compared to specialized vocational rehabilitation alone in individuals with an ABI.

## Complete Care Pathways



The goal in any rehabilitation stream is to provide seamless care from the onset of injury to the ultimate recovery. As this module has demonstrated, the continuum of ABI care involves acute interventions with a transition to some combination of rehabilitation therapies. This section aims to identify studies which have compared pathways of care combining several rehabilitation strategies.

**TABLE 11 | Comprehensive Care Pathways for Individuals with an ABI**

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<p><a href="#">Glintborg et al.</a> (2016) Denmark PCT N=82</p>	<p><b>Population:</b> ABI; <i>KORE group</i> (n=27): Mean Age=53.6yr; Gender: Male=16, Female=11; Diagnosis: Apoplexia=21, TBI=6. <i>ALT group</i> (n=18): Mean Age=52yr; Gender: Male=14, Female=4; Diagnosis: Apoplexia=16, TBI=2. <i>SR group</i> (n=37): Mean Age=53.4yr; Gender: Male=17, Female=20; Diagnosis: Apoplexia=25, Stroke=12.</p> <p><b>Intervention:</b> Participants who received the coordinated rehabilitation program (KORE) were compared with participants receiving standard treatment (SR group) and alternative treatment (ALT group).</p> <p><b>Outcome Measure:</b> Functional Independence Measure (FIM), Major Depression Inventory (MDI), WHO-Quality of Life-BREF (WHOQOL-BREF), Impact on Autonomy and Independence Questionnaire (IPAQQ-DK).</p>	<ol style="list-style-type: none"> <li>FIM improved significantly in all groups from hospitalization after ABI to discharge (KORE: p&lt;0.001, r=0.56; ALT: p&lt;0.001, r=0.62; SR: p&lt;0.001, r=0.58). At 1yr post discharge, only the ALT group had a significant increase in total FIM score (p&lt;0.001, r=0.48). There were significant differences between groups in total FIM score at discharge (p&lt;0.001). Post hoc analysis showed the ALT groups FIM score was significantly lower than the KORE group (p&lt;0.01, r=0.36).</li> <li>Signs of clinical depression at discharge were observed in 30% of clients in the KORE group and 22% of those in the ALT group. MDI scores increased non-significantly from discharge to 1-2yr post discharge. Depression rates did not differ significantly between groups at any time point.</li> <li>No significant change in any of the QOL scores from discharge to 1-2yr post discharge was recorded. In the KORE and ALT groups, 74% and 77% of clients respectively, reported being dissatisfied with their physical QOL.</li> <li>Indoor autonomy significantly improved from discharge to 1-2yr post discharge in both the KORE group (p&lt;0.001, r=0.43) and the ALT group (p&lt;0.001, r=0.49). The ALT group also reported significantly negative changes in family roles (p&lt;0.05, r=0.38) and a reduction in outdoor problems (p&lt;0.001, r=0.50).</li> </ol>
<p><a href="#">Andelic et al.</a> (2014) Norway Case Control N=59</p>	<p><b>Population:</b> TBI; <i>Continuous Group</i> (n=30): Gender: Male=23, Female=7. <i>Broken-Chain Group</i> (n=29): Gender: Male=22, Female=7.</p> <p><b>Intervention:</b> Two rehabilitation trajectories were explored: continuous and broken chain. Clinical data on patients who had been admitted to rehabilitation between 2005 and 2007, and had received 6wk, 1yr and 5yr follow-ups post-injury was analyzed.</p> <p><b>Outcome Measure:</b> Disability Rating Scale (DRS) and costs of treatment.</p>	<ol style="list-style-type: none"> <li>Patients in the continuous chain group experienced an additional 4.06 points gain in DRS compared to the Broken-chain group (19.40 versus 23.46).</li> <li>The cost for the continuous group from acute care through to rehabilitation was 37,000 NOK (approx. \$6,075.5 USD) less than the broken chain group.</li> </ol>

Author Year Country Research Design PEDro Sample Size	Methods	Outcome
<a href="#">Harradine et al.</a> (2004) Australia Cohort N=198	<p><b>Population:</b> TBI; <i>Urban (n=147)</i>: Mean Age=32.1yr; Gender: Male=117, Female=30. <i>Rural (n=51)</i>: Mean Age=32.1yr; Gender: Male=38, Female=13.</p> <p><b>Intervention:</b> Questionnaires were administered to patients at rehabilitation admission then again at 18mo follow-up. Patients were compared based on where they lived (urban or rural).</p> <p><b>Outcome Measure:</b> Disability Rating Scale (DRS), Mayo–Portland Adaptability Inventory (MPAI), General Health Questionnaire 28-item version (GHQ-28), and Medical Outcomes Short Form Health Survey (SF-36).</p>	<ol style="list-style-type: none"> <li>1. There were no significant differences between the two groups for scores on the DRS, MPAI, GHQ-28, and SF-36 questionnaires at 18mo follow-up.</li> <li>2. There were no significant differences between the two groups in return-to-work rate or functional outcomes at follow-up.</li> <li>3. In both groups, fewer patients were living alone compared to pre-injury; more were living with parents post injury, but this did not reach statistical significance.</li> </ol>
<a href="#">McLaughlin &amp; Peters</a> (1993) USA Cohort N=31	<p><b>Population:</b> ABI; <i>Step-Up Group (n=19)</i>: Mean Age=26.6yr; Gender: Male=18, Female=13; Mean Time Post injury=16.68 mo. <i>Inpatient Group (n=12)</i>: Mean Age=26.6yr, Gender: Male=18, Female=13; Mean Time Post Injury=18.3 mo.</p> <p><b>Intervention:</b> Patients in the Step-Up Group participated in a transitional living setting inpatient rehabilitation program and were compared with patients receiving inpatient rehabilitation alone. Data collected over 18mo through a post discharge survey.</p> <p><b>Outcome Measure:</b> Rancho Los Amigos Levels of Cognitive Functioning Scale (RLAS), Barthel Index (BI), and surveys on independent living and performance post-discharge.</p>	<ol style="list-style-type: none"> <li>1. Patients in the Step-Up Group reported greater functional independence on the BI than patients who received inpatient rehabilitation alone.</li> <li>2. Although patients in the Step-Up group reported better independent skills, they did not differ with the inpatient group on employment rates, participation in volunteer work or in RLAS scores.</li> </ol>

## Discussion

These studies re-affirm many of the concerns already noted in this module. There is significant heterogeneity in the care received by individuals with ABI and the direct comparison of complete systems is difficult. No matter what health care system is assessed, budgetary concerns play a role in the accessibility of care. As a result, difficult decisions need to be made regarding resource allocation. Despite financial concerns, Khan et al. (2002) provide encouraging news regarding decreases in LOS and fiscal savings brought on by an integrated ABI system in Canada. The authors state that care needs to be taken to ensure that savings do not arise from sacrifices in quality of care but rather from the improvement of systematic inefficiencies. Moreover, Andelic et al. (2014) report that a continuous chain of treatment and interventions worked out to be more cost-effective than the ‘broken chain’ format of rehabilitation with patients receiving inconsistent interventions. Thus, patients transitioning smoothly through the continuum of care not only benefit in terms of functional and cognitive gains, but approximately \$6,075.5 USD per patient was saved (Andelic et al., 2014). Finally, Harradine et al. (2004)



note that co-ordination of regional facilities resulted in an equal availability of resources despite geographic challenges in New South Wales, Australia.

Continuity and accessibility of services is crucial to allow patients the greatest opportunities for rehabilitation and recovery post ABI. Regional differences in resource availability need to be taken into consideration, along with patient demographics, so that the correct pathway decisions can be made. An alternative model of care is a comprehensive rehabilitation case management approach; this was implemented within a brain injury rehabilitation service and evaluated by Kennedy et al. (2012). A series of interviews with case managers and brain injury staff revealed that the new model provided a consistent and continuous transition through the rehabilitation continuum. Furthermore, Simpson et al. (2018) investigated the model of community-based case management in New South Wales, which was best characterized as a direct service model. No significant differences were found when comparing adult to paediatric programs. More time was spent on the task of support provision in rural as opposed to metropolitan sites. The services offered were rated as holistic, rather than service-led and decision making was routinely shared between staff and their clients.

### Conclusions

*There is level 2 evidence that individuals post ABI living in both rural and urban settings may have greater functional gains from an integrated network of inpatient, outpatient, and community services compared to standard inpatient rehabilitation.*



#### KEY POINTS

- Although continuity of care has been shown to be beneficial in optimizing recovery, there is insufficient evidence to draw conclusions regarding the ideal structure of a complete model of ABI care. Further research is required in determining the ideal structure of a complete model of ABI care.

## Conclusion

As stated previously, this module is not concerned with the individual effect of an intervention, but rather the comparative effect of different models of rehabilitation. Overall, the majority of the literature provides support for the effectiveness of earlier (versus late), and higher intensity (versus standard) inpatient or outpatient rehabilitation. In terms of guideline implementation, although guidelines themselves can foster improvement, there need to be specific strategic plans in place for their uptake in order to increase compliance and lead to improved patient outcomes. On an institutional level, there can be a variety of differences between institutions in terms of the provision of care, which can be due to differences in resources, staffing, staff training, and education.

The ultimate goal of any rehabilitation program is to fully reintegrate an individual back into society, both socially and vocationally. With this in mind, vocational rehabilitation programs which take a more



holistic approach have generally been found to be more effective. Further information on community reintegration can be found in Module 13.

Lastly, the issue of heterogeneous complete care pathways raises the concern that individuals with similar needs and functional status can undergo drastically different trajectories of rehabilitation care. Although no consensus currently exists for the optimal trajectory of care, institutions should prioritize needs assessments of their patients to determine the best course of treatment. Further research is needed in this area to help develop standardized trajectories across clinical settings.

## Reference List

Accreditation Canada. (2008). *Accreditation Canada*.

Andelic, N., Ye, J., Tornas, S., Roe, C., Lu, J., Bautz-Holter, E., Moger, T., Sigurdardottir, S., Schanke, A.-K., & Aas, E. (2014, 2014-12-02). Cost-effectiveness analysis of an early-initiated, continuous chain of rehabilitation after severe traumatic brain injury. *Journal of neurotrauma*, *31*(14), 1313-1320.

Ashley, M. J., Ashley, J., & Kreber, L. (2012). Remediation of information processing following traumatic brain injury: A community-based rehabilitation approach. *NeuroRehabilitation*, *31*(1), 31-39.

Badjatia, N., Carney, N., Croco, T., Fallat, M., Hennes, H., Jagoda, A., Jennigan, S., & Lerner, E. (2007). *Guidelines for the Pre-hospital Management of Severe Traumatic Brain Injury* (2 ed.). Brain Trauma Foundation.

Baethmann, A., Chapuis, D., & Wirth, A. (1999). System analysis of patient management during the pre- and early clinical phase in severe head injury. *Acta Neurochir Suppl*, *73*, 93-97.

Bender, A., Bauch, S., & Grill, E. (2014). Efficacy of a post-acute interval inpatient neurorehabilitation programme for severe brain injury. *Brain Injury*, *28*(1), 44-50.

Blackerby, W. (1990). Intensity of rehabilitation and length of stay. *Brain Injury*, *4*(2), 167-173.

Brain Injury Association of America. (2015). *Learn about the Brain Injury Association of America*.

Brain Injury Association of Canada. (2015). *Brain Injury Association of Canada (BIAC)*.

Braunling-McMorrow, D., Dollinger, S. J., Gould, M., Neumann, T., & Heiligenthal, R. (2010). Outcomes of post-acute rehabilitation for persons with brain injury. *Brain Injury*, *24*(7-8), 928-938.

Bulger, E. M., Nathens, A. B., Rivara, F. P., MacKenzie, E., Sabath, D. R., & Jurkovich, G. J. (2007). National variability in out-of-hospital treatment after traumatic injury. *Annals of emergency medicine*, *49*(3), 293-301.

Campbell, M. (2000). *Rehabilitation for traumatic brain injury: physical therapy practice in context* (2 ed.). Churchill Livingstone

Canadian Institute for Health Information. (2014). *National Rehabilitation Reporting System, Data Quality Documentation, 2013-2014*. CIHI.

Carney, N., Ghajar, J. (2007). Introduction. *Journal of Neurotrauma*, 24((supplement 1): S-1-S-2.).

Centers for Disease Control. (2008). *Centers for Disease Control (CDC)*.

Centers for Disease Control and Prevention. (2018a). *Surveillance of TBI-Related Emergency Department Visits, Hospitalizations, and Deaths*.

Centers for Disease Control and Prevention. (2018b). *TBI-related Emergency Department Visits, Hospitalizations, and Deaths (EDHDs)*.

Chan, L., Doctor, J., Temkin, N., MacLehose, R. F., Esselman, P., Bell, K., & Dikmen, S. (2001). Discharge disposition from acute care after traumatic brain injury: the effect of insurance type. *Archives of physical medicine and rehabilitation*, 82(9), 1151-1154.

Cicerone, K. D., Mott, T., Azulay, J., & Friel, J. C. (2004). Community integration and satisfaction with functioning after intensive cognitive rehabilitation for traumatic brain injury. *Archives of physical medicine and rehabilitation*, 85(6), 943-950.

Cifu, D. X., Kreutzer, J. S., Kolakowsky-Hayner, S. A., Marwitz, J. H., & Englander, J. (2003). The relationship between therapy intensity and rehabilitative outcomes after traumatic brain injury: A multicenter analysis. *Archives of physical medicine and rehabilitation*, 84(10), 1441-1448.

Citerio, G., Stocchetti, N., Cormio, M., Beretta, L., Galli, D., & Pesenti, A. (2003). Application of guidelines for severe head trauma: data from an Italian database. *European Journal of Emergency Medicine*, 10(1), 68-72.

Colantonio, A., Ratcliff, G., Chase, S., Kelsey, S., Escobar, M., & Vernich, L. (2004). Long term outcomes after moderate to severe traumatic brain injury. *Disability & Rehabilitation*, 26(5), 253-261.

Cope, D., & Hall, K. (1982). Head injury rehabilitation: benefit of early intervention. *Archives of physical medicine and rehabilitation*, 63(9), 433-437.

Cope, D. N. (1995). The effectiveness of traumatic brain injury rehabilitation: A review. *Brain Injury*, 9(7), 649-670.

Cope, D. N., Mayer, N. H., & Cervelli, L. (2005). Development of systems of care for persons with traumatic brain injury. *The Journal of head trauma rehabilitation*, 20(2), 128-142.

Cullen, N. (2007). Canadian healthcare perspective in traumatic brain injury rehabilitation. *The Journal of head trauma rehabilitation*, 22(4), 214-220.

- Cullen, N., Vimalasan, K., & Taggart, C. (2013). Efficacy of a functionally-based neurorehabilitation programme: A retrospective case-matched study of rehabilitation outcomes following traumatic brain injury. *Brain Injury*, 27(7-8), 799-806.
- Demetriades, D., Kimbrell, B., Salim, A., Velmahos, G., Rhee, P., Preston, C., Gruzinski, G., & Chan, L. (2005). Trauma Deaths in a Mature Urban Trauma System: Is “Trimodal” Distribution a Valid Concept? *Journal of the American College of Surgeons*, 201(3), 343-348.
- Drag, L., Renninger, C., King, R., & Hoblyn, J. (2013, Jan-Feb). Predictors of inpatient and outpatient healthcare utilization in veterans with traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 28(1), 39-47.
- DuBose, J. J., Browder, T., Inaba, K., Teixeira, P. G., Chan, L. S., & Demetriades, D. (2008). Effect of trauma center designation on outcome in patients with severe traumatic brain injury. *Archives of Surgery*, 143(12), 1213-1217.
- Edwards, M., McNeil, J., & Greenwood, R. (2003). Process and outcome during early inpatient rehabilitation after brain injury.
- Eicher, V., Murphy, M. P., Murphy, T. F., & Malec, J. F. (2012). Progress assessed with the Mayo-Portland Adaptability Inventory in 604 participants in 4 types of post-inpatient rehabilitation brain injury programs. *Archives of physical medicine and rehabilitation*, 93(1), 100-107.
- Espinosa-Aguilar, A., Reyes-Morales, H., Huerta-Posada, C. E., de León, I. L.-P., López-López, F., Mejía-Hernández, M., Mondragón-Martínez, M. A., Calderón-Téllez, L. M., Amezcua-Cuevas, R. L., & Rebollar-González, J. A. (2008). Design and validation of a critical pathway for hospital management of patients with severe traumatic brain injury. *Journal of Trauma-Injury, Infection, and Critical Care*, 64(5), 1327-1341.
- European Brain Injury Society. (2015). *European Brain Injury Society*.
- Evans, R. W. (1997). Postacute neurorehabilitation: roles and responsibilities within a national information system. *Archives of physical medicine and rehabilitation*, 78(8), S17-S25.
- Fakhry, S. M., Trask, A. L., Waller, M. A., & Watts, D. D. (2004). Management of brain-injured patients by an evidence-based medicine protocol improves outcomes and decreases hospital charges. *Journal of Trauma-Injury, Infection, and Critical Care*, 56(3), 492-500.
- Faul, M., Wald, M. M., Rutland-Brown, W., Sullivent, E. E., & Sattin, R. W. (2007). Using a cost-benefit analysis to estimate outcomes of a clinical treatment guideline: testing the Brain Trauma Foundation guidelines for the treatment of severe traumatic brain injury. *Journal of Trauma-Injury, Infection, and Critical Care*, 63(6), 1271-1278.

[Record #1 is using a reference type undefined in this output style.]

Formisano, R., Azicnuda, E., Sefid, M. K., Zampolini, M., Scarponi, F., & Avesani, R. (2017, 2016 Oct 01 2017-09-25). Early rehabilitation: Benefits in patients with severe acquired brain injury. *Neurological Sciences, 38*(1), 181-184.

Foster, M., & Tilse, C. (2003). Referral to rehabilitation following traumatic brain injury: a model for understanding inequities in access. *Social science & medicine, 56*(10), 2201-2210.

Gabriel, E. J., Ghajar, J., Jagoda, A., Pons, P. T., Scalea, T., Walters, B. C., & Brain, T. F. (2002). Guidelines for prehospital management of traumatic brain injury. *Journal of neurotrauma, 19*(1), 111.

Geeraerts, T., Velly, L., Abdenmour, L., Asehnoune, K., Audibert, G., Bouzat, P., Bruder, N., Carrillon, R., Cottenceau, V., Cotton, F., Courtil-Teyssedre, S., Dahyot-Fizelier, C., Dailler, F., David, J. S., Engrand, N., Fletcher, D., Francony, G., Gergelé, L., Ichai, C., Javouhey, E., Leblanc, P. E., Lieutaud, T., Meyer, P., Mirek, S., Orliaguet, G., Proust, F., Quintard, H., Ract, C., Srairi, M., Tazarourte, K., Vigué, B., & Payen, J. F. (2018). Management of severe traumatic brain injury (first 24 hours). *Anaesthesia critical care & pain medicine, 37*(2), 171-186.

Glintborg, C., & Hansen, T. G. (2016). Bio-psycho-social effects of a coordinated neurorehabilitation programme: A naturalistic mixed methods study. *NeuroRehabilitation, 38*(2), 99-113.

Godbolt, A. K., Stenberg, M., Jakobsson, J., Sorjonen, K., Krakau, K., Stalnacke, B. M., & Nygren DeBoussard, C. (2015, May 3). Subacute complications during recovery from severe traumatic brain injury: frequency and associations with outcome. *BMJ Open, 5*(4), e007208.

Griesdale, D. E., Örténwall, V., Norena, M., Wong, H., Sekhon, M. S., Kolmodin, L., Henderson, W. R., & Dodek, P. (2015). Adherence to guidelines for management of cerebral perfusion pressure and outcome in patients who have severe traumatic brain injury. *Journal of critical care, 30*(1), 111-115.

Harradine, P. G., Winstanley, J. B., Tate, R., Cameron, I. D., Baguley, I. J., & Harris, R. D. (2004). Severe traumatic brain injury in New South Wales: comparable outcomes for rural and urban residents. *Medical journal of Australia, 181*(3), 130-135.

Harrington, D. T., Connolly, M., Biffi, W. L., Majercik, S. D., & Cioffi, W. G. (2005). Transfer times to definitive care facilities are too long: a consequence of an immature trauma system. *Annals of surgery, 241*(6), 961.

Harris, O. A., Bruce, C. A., Reid, M., Cheeks, R., Easley, K., Surles, M. C., Pan, Y., Rhoden-Salmon, D., Webster, D., & Crandon, I. (2008). Examination of the management of traumatic brain injury in

the developing and developed world: focus on resource utilization, protocols, and practices that alter outcome.

- Hart, T., Whyte, J., Poulsen, I., Kristensen, K. S., Nordenbo, A. M., Chervoneva, I., & Vaccaro, M. J. (2016, Dec). How Do Intensity and Duration of Rehabilitation Services Affect Outcomes From Severe Traumatic Brain Injury? A Natural Experiment Comparing Health Care Delivery Systems in 2 Developed Nations. *Arch Phys Med Rehabil*, *97*(12), 2045-2053.
- Hassett, L. M., Moseley, A. M., Tate, R. L., Harmer, A. R., Fairbairn, T. J., & Leung, J. (2009). Efficacy of a fitness centre-based exercise programme compared with a home-based exercise programme in traumatic brain injury: a randomized controlled trial. *Journal of Rehabilitation Medicine*, *41*(4), 247-255.
- Heinemann AW, S. V., Cichowski K, Tuel SM, Betts HB. (1990). Functional outcome following traumatic brain injury rehabilitation: Translating research into policy. *Journal of Neurologic Rehabilitation*, *4*, 27-37.
- Hesdorffer, D. C., & Ghajar, J. (2007, Oct). Marked improvement in adherence to traumatic brain injury guidelines in United States trauma centers. *J Trauma*, *63*(4), 841-847; discussion 847-848.
- High Jr, W. M., Roebuck-Spencer, T., Sander, A. M., Struchen, M. A., & Sherer, M. (2006). Early versus later admission to postacute rehabilitation: Impact on functional outcome after traumatic brain injury. *Archives of physical medicine and rehabilitation*, *87*(3), 334-342.
- Hyder, A. A., Wunderlich, C. A., Puvanachandra, P., Gururaj, G., & Kobusingye, O. C. (2007). The impact of traumatic brain injuries: a global perspective. *NeuroRehabilitation*, *22*(5), 341-353.
- International Brain Injury Association. (2008). *International Brain Injury Association*
- Jeyaraj, J. A., Clendenning, A., Bellemare-Lapierre, V., Iqbal, S., Lemoine, M.-C., Edwards, D., & Korner-Bitensky, N. (2013, 2014-12-02). Clinicians' perceptions of factors contributing to complexity and intensity of care of outpatients with traumatic brain injury. *Brain Injury*, *27*(12), 1338-1347.
- Juelsgaard, J., Rognås, L., Knudsen, L., Hansen, T. M., & Rasmussen, M. (2018). Prehospital treatment of patients with acute intracranial pathology - adherence to guidelines and blood pressure recommendations by the Danish Air Ambulance. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine*, *26*(1), 68-68.
- Kennedy, N., Barnes, J., Rose, A., & Veitch, C. (2012). Clinicians' expectations and early experiences of a new comprehensive rehabilitation case management model in a specialist brain injury rehabilitation unit. *Brain Impairment*, *13*(1), 62-71.

- Smith, J., Smith, D., No map, no guide: Family caregivers' perspectives on their journeys through the system (2000) *Case Management Journals*, 2 (2), pp. 27-33
- Turner, B., Fleming, J., Cornwell, P., Worrall, L., Ownsworth, T., Haines, T., Kendall, M., Chenoweth, L., A qualitative study of the transition from hospital to home for individuals with acquired brain injury and their family caregivers (2007) *Brain Injury*, 21 (11), pp. 1119-1130.
- Turner, B., Fleming, J., Ownsworth, T., Cornwell, P., The transition from hospital to home for individuals with acquired brain injury: A literature review and research recommendations (2008) *Disability and Rehabilitation*, 30 (16), pp. 1153-1176
- Turner, B., Ownsworth, T., Cornwell, P., Fleming, J., Reengagement in meaningful occupations during the transition from hospital to home for people with acquired brain injury and their family caregivers (2009) *American Journal of Occupational Therapy*, 63 (5), pp. 609-620
- Turner, B., Fleming, J., Cornwell, P., Haines, T., Ownsworth, T., Profiling early outcomes during the transition from hospital to home after brain injury (2009) *Brain Injury*, 23 (1), pp. 551-560
- Turner, B., Fleming, J., Ownsworth, T., Cornwell, P., Perceptions of recovery during the early transition phase from hospital to home following acquired brain injury: A journey of discovery (2011) *Neuropsychological Rehabilitation*, 21 (1), pp. 64-91
- Voss, R., Gardner, R., Baier, R., Butterfield, K., Lehrman, S., Gravenstein, S., The care transitions intervention: Translating from efficacy to effectiveness (2011) *Archives of Internal Medicine*, 171 (14), pp. 1232-1237
- Kesinger, M. R., Nagy, L. R., Sequeira, D. J., Charry, J. D., Puyana, J. C., & Rubiano, A. M. (2014). A standardized trauma care protocol decreased in-hospital mortality of patients with severe traumatic brain injury at a teaching hospital in a middle-income country. *Injury*, 45(9), 1350-1354.
- Khan, S., Khan, A., & Feyz, M. (2002). Decreased length of stay, cost savings and descriptive findings of enhanced patient care resulting from an integrated traumatic brain injury programme. *Brain Injury*, 16(6), 537-554.
- Khormi, Y. H., Gosadi, I., Campbell, S., Senthilselvan, A., O'Kelly, C., & Zygun, D. (2018). Adherence to Brain Trauma Foundation Guidelines for Management of Traumatic Brain Injury Patients and Its Effect on Outcomes: Systematic Review. *Journal of neurotrauma*, 35(13), 1407-1418.
- Kim, H., Colantonio, A., Deber, R., & Vernich, L. (2006). Discharge destination from acute care after traumatic brain injury. *The Canadian journal of neurological sciences*, 33(1), 48-52.



- Kramer, A. H., & Zygun, D. A. (2013). Declining mortality in neurocritical care patients: A cohort study in Southern Alberta over eleven years. *Canadian Journal of Anesthesia*, *60*(10), 966-975.
- Lamontagne, M. E., Bayley, M. T., Marshall, S., Kua, A., Marier-Deschenes, P., Allaire, A. S., Schmouth, M. E., Kagan, C., Truchon, C., & Swaine, B. (2018). Assessment of users' needs and expectations toward clinical practice guidelines to support the rehabilitation of adults with moderate to severe traumatic brain injury. *Journal of Head Trauma Rehabilitation*, *33*(5), 288-295.
- Lamontagne, M. E., Gargaro, J., Marier-Deschenes, P., Truchon, C., Bayley, M. T., Marshall, S., Kagan, C., Briere, A., & Swaine, B. (2018). A survey of perceived implementation gaps for a clinical practice guideline for the rehabilitation of adults with moderate to severe traumatic brain injury. *Journal of Head Trauma Rehabilitation*, *33*(5), 306-316.
- Leith, K. H., Phillips, L., & Sample, P. L. (2004). Exploring the service needs and experiences of persons with TBI and their families: the South Carolina experience. *Brain Injury*, *18*(12), 1191-1208.
- León-Carrión, J., MacHuca-Murga, F., Solís-Marcos, I., León-Domínguez, U., & Domínguez-Morales, M. D. R. (2013). The sooner patients begin neurorehabilitation, the better their functional outcome. *Brain Injury*, *27*(10), 1119-1123.
- Maas, A., Dearden, M., Teasdale, G., Braakman, R., Cohadon, F., Iannotti, F., Karimi, A., Lapierre, F., Murray, G., & Ohman, J. (1997). EBIC-guidelines for management of severe head injury in adults. *Acta neurochirurgica*, *139*(4), 286-294.
- Mackay, L., Bernstein, B., Chapman, P., Morgan, A., & Milazzo, L. (1992). Early intervention in severe head injury: long-term benefits of a formalized program. *Archives of physical medicine and rehabilitation*, *73*(7), 635-641.
- MacKenzie, E. J., Rivara, F. P., Jurkovich, G. J., Nathens, A. B., Frey, K. P., Egleston, B. L., Salkever, D. S., Weir, S., & Scharfstein, D. O. (2007). The national study on costs and outcomes of trauma. *Journal of Trauma-Injury, Infection, and Critical Care*, *63*(6), S54-S67.
- Malec, J. F., & Degiorgio, L. (2002). Characteristics of successful and unsuccessful completers of 3 postacute brain injury rehabilitation pathways. *Archives of physical medicine and rehabilitation*, *83*(12), 1759-1764.
- March, E., Salzberg, M., & Hopwood, M. (2013). Acquired brain injury needs to be acknowledged at the national mental health policy level. *Australian and New Zealand Journal of Psychiatry*, *47*(3), 212-216.
- Martelli, M. F., Zasler, N. D., & Tiernan, P. (2012). Community based rehabilitation: Special issues. *NeuroRehabilitation*, *31*(1), 3-18.

- Matta, B., & Menon, D. (1996). Severe head injury in the United Kingdom and Ireland: A survey of practice and implications for management. *Critical care medicine*, 24(10), 1743-1748.
- Mazaux, J., & Richer, E. (1998). Rehabilitation after traumatic brain injury in adults. *Disability & Rehabilitation*, 20(12), 435-447.
- McLaughlin, A. M., & Peters, S. (1993). Evaluation of an innovative cost-effective programme for brain injury patients: response to a need for flexible treatment planning. *Brain Injury*, 7(1), 71-75.
- Moseley, A. M., Herbert, R. D., Sherrington, C., & Maher, C. G. (2002). Evidence for physiotherapy practice: a survey of the Physiotherapy Evidence Database (PEDro). *Aust J Physiother*, 48(1), 43-49.
- Myburgh, J. A., Cooper, D. J., Finfer, S. R., Venkatesh, B., Jones, D., Higgins, A., Bishop, N., Higlett, T., & Australian, A. T. B. I. S. I. f. t. (2008). Epidemiology and 12-month outcomes from traumatic brain injury in Australia and New Zealand. *Journal of Trauma-Injury, Infection, and Critical Care*, 64(4), 854-862.
- National Association of State Head Injury Administrators. (2008 ). *National Association of State Head Injury Administrators*.
- National Data and Statistical Center. (2008). *The Traumatic Brain Injury Model Systems of Care*
- National Institute of Health. (1998). *Rehabilitation of persons with traumatic brain injury*. NIH.
- New Zealand Guidelines Group. (2006). *Traumatic Brain Injury: Diagnosis, Acute Management and Rehabilitation*
- Olesen, J., & Freund, T. F. (2006). European Brain Council: partnership to promote European and national brain research. *Trends in Neurosciences*, 29(9), 493-495.
- Owensworth, T., Fleming, J., Shum, D., Kuipers, P., & Strong, J. (2008). Comparison of individual, group and combined intervention formats in a randomized controlled trial for facilitating goal attainment and improving psychosocial function following acquired brain injury. *Journal of Rehabilitation Medicine*, 40(2), 81-88.
- Palmer, S., Bader, M. K., Qureshi, A., Palmer, J., Shaver, T., Borzatta, M., & Stalcup, C. (2001). The impact on outcomes in a community hospital setting of using the AANS traumatic brain injury guidelines. *Journal of Trauma and Acute Care Surgery*, 50(4), 657-664.

- Peirone, E., Gorla, P. F., & Anselmino, A. (2014, 2014-12-02). A dual-task home-based rehabilitation programme for improving balance control in patients with acquired brain injury: A single-blind, randomized controlled pilot study. *Clinical Rehabilitation*, *28*(4), 329-338.
- Poncet, F., Swaine, B., Migeot, H., Lamoureux, J., Picq, C., & Pradat, P. (2018, Jun). Effectiveness of a multidisciplinary rehabilitation program for persons with acquired brain injury and executive dysfunction. *Disabil Rehabil*, *40*(13), 1569-1583.
- Ponsford, J., Harrington, H., Olver, J., & Roper, M. (2006). Evaluation of a community-based model of rehabilitation following traumatic brain injury. *Neuropsychological Rehabilitation*, *16*(3), 315-328.
- Powell, J., Heslin, J., & Greenwood, R. (2002). Community based rehabilitation after severe traumatic brain injury: a randomised controlled trial. *Journal of Neurology, Neurosurgery & Psychiatry*, *72*(2), 193-202.
- Rapport, L. J., Bryer, R. C., & Hanks, R. A. (2008). Driving and Community Integration After Traumatic Brain Injury. *Archives of physical medicine and rehabilitation*, *89*(5), 922-930.
- Reid-Arndt, S. A., Frank, R. G., & Hagglund, K. J. (2010). Brain injury and health policy: Twenty-five years of progress. *Journal of Head Trauma Rehabilitation*, *25*(2), 137-144.
- Rosenbaum, A. M., Gordon, W. A., Joannou, A., & Berman, B. A. (2018). Functional outcomes following post-acute rehabilitation for moderate-to-severe traumatic brain injury. *Brain Injury*, *32*(7), 907-914.
- Saatman, K. E., Duhaime, A.-C., Bullock, R., Maas, A. I., Valadka, A., & Manley, G. T. (2008). Classification of traumatic brain injury for targeted therapies. *Journal of neurotrauma*, *25*(7), 719-738.
- Sackett DL, S. S., Richardson WS, Rosenberg W, Hayes RB. (2000). *Evidence-based medicine: how to practice and teach EBM* (2nd ed. ed.).
- Saherwala, A. A., Bader, M. K., Stutzman, S. E., Figueroa, S. A., Ghajar, J., Gorman, A. R., Minhajuddin, A., & Olson, D. M. (2018). Increasing Adherence to Brain Trauma Foundation Guidelines for Hospital Care of Patients With Traumatic Brain Injury. *Critical care nurse*, *38*(1), e11-e20.
- Sample, P. L., & Langlois, J. A. (2005). Linking people with traumatic brain injury to services: successes and challenges in Colorado. *The Journal of head trauma rehabilitation*, *20*(3), 270-278.
- Semlyen, J. K., Summers, S. J., & Barnes, M. P. (1998). Traumatic brain injury: efficacy of multidisciplinary rehabilitation. *Archives of physical medicine and rehabilitation*, *79*(6), 678-683.

- Shafi, S., Barnes, S. A., Millar, D., Sobrino, J., Kudyakov, R., Berryman, C., Rayan, N., Dubiel, R., Coimbra, R., Magnotti, L. J., Vercruyssen, G., Scherer, L. A., Jurkovich, G. J., & Nirula, R. (2014). Suboptimal compliance with evidence-based guidelines in patients with traumatic brain injuries: Clinical article. *Journal of Neurosurgery*, *120*(3), 773-777.
- Shiel, A., Burn, J. P., Henry, D., Clark, J., Wilson, B., Burnett, M., & McLellan, D. (2001). The effects of increased rehabilitation therapy after brain injury: results of a prospective controlled trial. *Clinical Rehabilitation*, *15*(5), 501-514.
- Shigemori, M., & Tokutomi, T. (2002). Result of nationwide survey of the management of severe head injury in Japan. *Neurological research*, *24*(1), 41-44.
- Simpson, G., Mitsch, V., Doyle, M., Forman, M., Young, D., Solomon, V., MacPherson, M., Gillett, L., & Strettles, B. (2018). Investigating the Model of Community-Based Case Management in the New South Wales Brain Injury Rehabilitation Program: A Prospective Multicenter Study. *The Journal of head trauma rehabilitation*, *33*(6), E38-E48.
- Sorinola, A., Buki, A., Sandor, J., & Czeiter, E. (2018). Effectiveness of Traumatic Brain Injury Management Guideline Introduction in Hungary. *Turk Neurosurg*, *28*(3), 410-415.
- Spivack, G., Spettell, C. M., Ellis, D. W., & Ross, S. E. (1992). Effects of intensity of treatment and length of stay on rehabilitation outcomes. *Brain Injury*, *6*(5), 419-434.
- Surgeons, A. C. o. (1999). *Committee on Trauma of the American College of Surgeons*.
- Talving, P., Karamanos, E., Teixeira, P. G., Skiada, D., Lam, L., Belzberg, H., Inaba, K., & Demetriades, D. (2013). Intracranial pressure monitoring in severe head injury: Compliance with Brain Trauma Foundation guidelines and effect on outcomes: A prospective study. *Journal of Neurosurgery*, *119*(5), 1248-1254.
- Talving, P., Karamanos, E., Teixeira, P. G., Skiada, D., Lam, L., Belzberg, H., Inaba, K., & Demetriades, D. (2013). Intracranial pressure monitoring in severe head injury: compliance with Brain Trauma Foundation guidelines and effect on outcomes: a prospective study: Clinical article. *Journal of Neurosurgery*, *119*(5), 1248-1254.
- Tarapore, P. E., Vassar, M. J., Cooper, S., Lay, T., Galletly, J., Manley, G. T., & Huang, M. C. (2016, 2016 Nov 15 2017-09-25). Establishing a traumatic brain injury program of care: Benchmarking outcomes after institutional adoption of evidence-based guidelines. *Journal of neurotrauma*, *33*(22), 2026-2033.

- Teasdale, G., Braakman, R., Cohadon, F., Dearden, M., Iannotti, F., Karimi, A., Lapierre, F., Maas, A., Murray, G., & Ohman, J. (1997). The European brain injury consortium. *Acta neurochirurgica*, *139*(9), 797-803.
- Tuel, S. M., Presty, S. K., Meythaler, J. M., Heinemann, A. W., & Katz, R. T. (1992). Functional improvement in severe head injury after readmission for rehabilitation. *Brain Injury*, *6*(4), 363-372.
- Wagner, A. K., Fabio, T., Zafonte, R. D., Goldberg, G., Marion, D. W., & Peitzman, A. B. (2003). Physical medicine and rehabilitation consultation: relationships with acute functional outcome, length of stay, and discharge planning after traumatic brain injury. *American journal of physical medicine & rehabilitation*, *82*(7), 526-536.
- Webb, P. M., & Glueckauf, R. L. (1994). The effects of direct involvement in goal setting on rehabilitation outcome for persons with traumatic brain injuries. *Rehabilitation Psychology*, *39*(3), 179.
- Willer, B., Button, J., & Rempel, R. (1999). Residential and home-based postacute rehabilitation of individuals with traumatic brain injury: a case control study. *Archives of physical medicine and rehabilitation*, *80*(4), 399-406.
- Zasler, N., Katz, D., & Zafonte, R. D. (2007). *Brain injury medicine: Principles and practice*. Demos Medical Publishing.
- Zhu, X. L., Poon, W. S., Chan, C. H., & Chan, S. H. (2001). Does intensive rehabilitation improve the functional outcome of patients with traumatic brain injury? Interim result of a randomized controlled trial. *British Journal of Neurosurgery*, *15*(6), 464-473.
- Zygun, D. A., Laupland, K. B., Hader, W. J., Kortbeek, J. B., Findlay, C., Doig, C. J., & Hameed, S. M. (2005). Severe traumatic brain injury in a large Canadian health region. *The Canadian journal of neurological sciences*, *32*(1), 87-92.